

Quantum geometry from phase space reduction

Florian Conrady* and Laurent Freidel†

Perimeter Institute for Theoretical Physics, Waterloo, Ontario, Canada

In this work we give an explicit isomorphism between the usual spin network basis and the direct quantization of the reduced phase space of tetrahedra. The main outcome is a formula that describes the space of $SU(2)$ invariant states by an integral over coherent states satisfying the closure constraint exactly, or equivalently, as an integral over the space of classical tetrahedra. This provides an explicit realization of theorems by Guillemin–Sternberg and Hall that describe the commutation of quantization and reduction. In the final part of the paper, we use our result to express the FK spin foam model as an integral over classical tetrahedra and the asymptotics of the vertex amplitude is determined.

I. INTRODUCTION

Recently a wealth of important developments has occurred following the proposal of new spin foam models for gravity [1, 2, 3, 4]. These models allow the inclusion of a non-trivial Immirzi parameter and have been shown to satisfy¹ two very important consistency requirements which have been eluding us for a long time: firstly, in the semiclassical limit they are asymptotically equivalent to the usual Regge discretisation of gravity, independently of the complexity of the underlying cell complex [5]. Secondly, they have been shown to possess $SU(2)$ spin network states as boundary states [6, 7].

We can link these new developments with the evolution of our understanding of the so-called simplicity constraints. These simplicity constraints state (in any dimension [8]) that the bivector used to contract the curvature tensor in the Einstein action comes from a frame field. In a simplicial context this constraint splits into three classes: there are the face simplicity constraints (among bivectors associated to the same face), the cross-simplicity constraints (among bivectors associated to faces of the same tetrahedron) and the volume constraints (among opposite bivectors). All these constraints are quadratic, but the volume constraint, because it depends on different tetrahedra, also involves the connection and is therefore extremely hard to quantise. The first major advance came from the work of Baez, Barrett and Crane [9, 10] who showed that it is possible to linearise the volume constraint and replace it by a constraint living only at one tetrahedron. This new constraint is the closure constraint and the discrete analogue of the Gauss law generating gauge transformations. The second key insight in this direction came from the work of Engle, Pereira and Rovelli [1] who showed that one can linearise the simplicity and cross simplicity constraints, opening the path toward a new way of constructing spin foam models and allowing the incorporation of the Immirzi parameter.

*Electronic address: fconrady@perimeterinstitute.ca

†Electronic address: lfreidel@perimeterinstitute.ca

¹ at least in the Riemannian sector

There is, however, one unsatisfactory point in this description: if one looks at a path integral version of these models [7] one quickly recognizes that the simplicity and cross simplicity constraints are imposed strongly, whereas the closure constraint is only imposed weakly as an expression of gauge invariance and becomes true only in the semiclassical limit. Such a distinction does not occur in the continuum theory where all simplicity constraints are imposed strongly [11]. This is related to the fact that the theory admits a geometrical interpretation (in terms of metrics) only in the semiclassical regime, which raised recent criticism [12]. The purpose of our work is to solve this problem and to show that the new spin foam model can, in fact, be written explicitly in terms of a sum of amplitudes, where all simplicity constraints are treated on the same footing, i.e. imposed strongly in the path integral, and where all the boundary spin networks have a geometrical interpretation even before taking the semiclassical limit.

It is also interesting to note that along with the construction of new models we are also witnessing a merging of two lines of thoughts on spin foam models that developed in parallel for a long time and are now finally intersecting. One line of thought, which is more canonical in spirit, can be traced back to the seminal work of Barbieri [13], who realized that spin network states of loop quantum gravity can be understood (and heuristically derived) by applying the quantization procedure to a collection of geometric tetrahedra in 3 spatial dimensions. This important work suggested that spin network states and quantum gravity may be about quantizing geometric structures. This idea was then quickly applied to the problem of quantization of a geometric simplex in $\mathbb{R}^4$, with a breakthrough result being the construction of the now famous Barrett–Crane model [9]. This is also the line of thought developed by Engle, Livine, Pereira, Rovelli in [4]. The main problem in this approach is that the geometry associated with the quantum tetrahedra is fuzzy and cannot be resolved sharply due to the noncommutativity of the geometric “flux” operators. This problem has haunted the field for a long time, since it prevents a priori a sharp coupling between neighboring 4d building blocks, and that was the main issue with the Barrett–Crane model.

The second line of thought can be traced back to the work of Reisenberger [14] who proposed to think about quantum gravity directly in terms of a path integral approach in which we integrate over classical configurations living on a 2–dimensional spine and where spin foam models arise from a type of bulk discretization. This line of reasoning is deeply related to the Plebanski reformulation of gravity as a constraint $SO(4)$ BF theory [15, 16]. In this approach it is quite easy to get a consistent gluing condition, however, it is much less trivial to find models which have a natural and simple algebraic expression in terms of recoupling coefficients and it is even more of a challenge to get $SU(2)$ boundary spin networks as boundary states.

The merging between these two approaches starts with a work of Livine and Speziale [17] who proposed to label spin network states not with the usual intertwiner basis, but with “coherent intertwiners” that are labelled by four vectors whose norm is fixed to be the area of the faces of the tetrahedron. This is almost the missing link between the two approaches and it has led to a very efficient and geometrical way of deriving the new spin foam models [2, 3].

There is, however, an important caveat. The coherent intertwiner resolves the fuzziness of the quantum space of a tetrahedron by construction, but it does so at a high price: the classical configuration no longer satisfies the closure constraint unless one takes the semiclassical limit and it is therefore not geometrical.

The resolution of these problems that we propose here is very simple in spirit: one should

use coherent states that are associated to the geometrical configuration (see [18] for an early attempt) and relate these “geometrical” states to the usual spin network states or the coherent intertwiner states.

In this paper, we give a construction of such states, we show that they form an over-complete basis of the space of four-valent intertwiners and we give the explicit isomorphism between this new basis of states, where the closure constraint is imposed strongly, and the coherent intertwiner basis.

The proof of this isomorphism amounts essentially to showing that “quantisation commutes with reduction” and utilises heavily the beautiful and seminal work of Guillemin and Sternberg [19]. All our proofs and derivations are self-contained and we do not assume any knowledge of the mathematical literature [20, 21, 22, 23, 38] on geometrical quantisation which is usually the arena in which this work is presented.

The main formula of the paper is described in section V. It expresses the decomposition of identity in the space of four-valent intertwiners $H_{\vec{j}} = (V^{j_1} \otimes \dots \otimes V^{j_4})^{\text{SU}(2)}$ as the following integral over coherent states satisfying the closure constraint, and hence as an integral over the space of classical tetrahedra:

$$\mathbb{1}_{\vec{j}} = \prod_i d_{j_i} \int_{P_{\vec{j}}^{(0)}} \prod_i d^2 N_i \delta \left(\sum_i j_i N_i \right) \rho_{\vec{j}}(N_i) ||\vec{j}, N_i\rangle \langle \vec{j}, N_i|| \quad (1)$$

$||\vec{j}, N_i\rangle$ are the invariant intertwiners of Livine & Speziale [17]. The delta function imposes the closure constraint strongly, so the integral is really an integral over the classical phase space. Finally, $\rho_{\vec{j}}(N_i)$ is a positive $\text{SU}(2)$ invariant function which will be explicitly determined in eq. (103) and which characterises the isomorphism between the usual spin network basis and this “classical” basis.

Once this identity is obtained we apply it to the spin foam models given in [2]: it allows us to formulate the path integral in terms of variables on which the closure constraints are imposed strongly. These variables are well-suited to analyze the asymptotic large spin behaviour of amplitudes, and we use them here to derive the asymptotics of the FK vertex amplitude. Shortly after this work a paper by Barrett et al. appeared in which the asymptotics of the vertex amplitude is derived as well [25].

Our paper is organized as follows: in section II we describe in detail the phase space of the classical tetrahedron and show that it can be obtained as a symplectic quotient. We also state the classical part of the Guillemin–Sternberg isomorphism, which is a remarkable isomorphism between a constrained phase space divided by the action of the gauge group and the *unconstrained* phase space divided by the complexification of the group. Section III provides a detailed discussion of coherent states and their link with geometrical quantisation. In section IV and V we present a “pedestrian” proof of the Guillemin–Sternberg theorem for tetrahedra and construct explicitly the isomorphism between the new geometrical basis and the coherent intertwiner basis. In the last section, we rewrite the $\text{FK}\gamma$ spin foam model in terms of the new tetrahedral states and exploit this to derive the asymptotics of the vertex amplitude.

II. ON THE SPACE OF SHAPE OF TETRAHEDRA

A. The classical phase space of the polygon

Given a four-tuple $\vec{j} \equiv (j_1, \dots, j_4)$ of positive real numbers j_i we consider the Polygon space $\mathcal{P}_{\vec{j}}$ which is the space of 4-sided polygons in $\mathbb{R}^3$ with edge length equal to j_i modulo rotations. This space is non-empty if and only if

$$2j_i \leq j_1 + \dots + j_4 \quad i = 1, \dots, 4. \quad (2)$$

It can be written as the symplectic quotient of $P_{\vec{j}} \equiv S_{j_1}^2 \times \dots \times S_{j_4}^2$ by $SU(2)$, where S_j^2 is the 2-sphere of radius j and $SU(2)$ acts on it by rotation. In this paper, we identify $\mathbb{R}^3$ with $\mathfrak{su}(2)^*$, we represent $\mathfrak{su}(2)$ as the space of anti-hermitian, traceless 2 by 2 matrices, and we identify $\mathfrak{su}(2)$ and its dual using the metric $(X|Y) \equiv -2\text{tr}(XY)$. In this description S_j^2 is a (co)adjoint orbit of $\mathfrak{su}(2)$:

$$S_j^2 = \{X = j(n\tau_3n^{-1}) \in \mathfrak{su}(2) \mid n \in SU(2)\}, \quad (3)$$

where $\tau_i \equiv -i\sigma_i/2$ and σ_i denote the Pauli matrices (note that the scalar product is such that $(\tau_a|\tau_b) = \delta_{ab}$). We use a notation where $X = jN$ denotes a point on S_j^2 and $N \equiv n\tau_3n^{-1}$ denotes the corresponding unit vector.

S_j^2 is a symplectic manifold and the associated Poisson structure² is given by

$$\{f, g\}(X) = \epsilon^{abc} X_a \partial_b f \partial_c g \quad (4)$$

with $X_b = (\tau_b|X)$. From this it is obvious that linear functions $h_Y(X) \equiv (Y|X)$ generate the $SU(2)$ action on S^2 . We denote by $\hat{Y}$ the corresponding vector fields:

$$\{h_Y, f\} = \hat{Y}f, \quad \text{with} \quad \hat{Y}f(X) = \left. \frac{d}{dt} f(e^{-tY} X e^{tY}) \right|_{t=0}. \quad (5)$$

The symplectic form on S_j^2 is given by

$$\omega_j(\hat{Y}_1, \hat{Y}_2) = h_{[Y_1, Y_2]} = \{h_{Y_1}, h_{Y_2}\}. \quad (6)$$

The first important feature of the Polygon space $\mathcal{P}_{\vec{j}}$ is that it can be written as a symplectic quotient. Namely, let us first define on $P_{\vec{j}} = S_{j_1}^2 \times \dots \times S_{j_4}^2$ the Lie algebra valued function

$$H(X_i) \equiv \sum_i X_i = \sum_i j_i N_i, \quad \text{and} \quad H_Y(X_i) \equiv (Y|H) = \sum_i h_Y(X_i). \quad (7)$$

² Given a Poisson bracket we can assign to any function h on phase space a Hamiltonian vector field $\hat{X}_h \equiv \{h, \cdot\}$. When this Poisson bracket is non degenerate it defines a symplectic form ω by

$$i_{\hat{X}_h} \omega = -dh, \quad \text{hence} \quad \omega(\hat{X}_{h_1}, \hat{X}_{h_2}) = \{h_1, h_2\}.$$

H_X is the Hamiltonian³ generating the diagonal action $e^X \in \text{SU}(2)$ on $P_{\vec{j}}$. We have the isomorphism

$$\mathcal{P}_{\vec{j}} = P_{\vec{j}}^{(0)} / \text{SU}(2), \quad \text{with} \quad P_{\vec{j}}^{(0)} \equiv H^{-1}(0). \quad (8)$$

This quotient is non singular iff $j_1 \pm j_2 \pm j_3 \pm j_4 \neq 0$ for any choice of sign. When this inequality is satisfied, the action of $\text{SO}(3)$ on $H^{-1}(0)$ possesses no fixed point and the quotient is a Hausdorff manifold [27].

It follows from this isomorphism that $\mathcal{P}_{\vec{j}}$ is a 2-dimensional symplectic manifold⁴. Symplectic coordinates can be explicitly constructed [27, 28, 29]: one defines $j_{12} \equiv |X_1 + X_2|$, the length of one diagonal of the 4-gon, and $\phi_{12} \in [0, 2\pi]$ as the angle between the two oriented triangles $(X_1, X_2, X_1 + X_2)$ and $(X_3, X_4, X_3 + X_4)$ which meet along this diagonal. Then,

$$\Omega = dj_{12} \wedge d\phi_{12}. \quad (9)$$

To show this, one computes the Hamiltonian flow generated by j_{12} , which is the “bending” flow generating rotations around the diagonal, i.e.

$$\{j_{12}, f\}(X) = \left. \frac{d}{dt} f(e^{-tN_{12}} X e^{tN_{12}}) \right|_{t=0}, \quad \text{with} \quad N_{12} \equiv \frac{X_1 + X_2}{|X_1 + X_2|}. \quad (10)$$

This result implies that $\{j_{12}, \phi_{12}\} = 1$, hence the form of the symplectic structure.

The next step is to demonstrate that the polygon space is a complex Kähler manifold and we first review some aspects of complex geometry on the sphere. The complex structure on the sphere is a map J such that $J(\partial_z) = i\partial_z$, $J(\partial_{\bar{z}}) = -i\partial_{\bar{z}}$, where the complex coordinates z are defined by stereographic projection⁵:

$$N(z) = n(z)\tau_3 n(z)^{-1}, \quad n(z) \equiv \frac{1}{\sqrt{1+|z|^2}} \begin{pmatrix} 1 & z \\ -\bar{z} & 1 \end{pmatrix}. \quad (12)$$

The symplectic structure on the sphere S_j^2 can be expressed in terms of these coordinates as

$$\omega_j = \frac{2j}{i} \frac{dz \wedge d\bar{z}}{(1+|z|^2)^2}, \quad (13)$$

its normalisation being such that $\int \omega_j = -4\pi j$. We can also determine the explicit expressions of the Hamiltonians $h_a(z) \equiv h_{\tau_a}(X(z))$ and the corresponding vector fields⁶ $\widehat{X}_a$ in

³ These functions are referred to as the “moment map” in the mathematical literature [20, 26], whereas in physics they are called a system of first class constraints.

⁴ There is a unique symplectic structure Ω on $P_{\vec{j}}^{(0)} / \text{SU}(2)$ such that $i^*(\sum_i \omega_{j_i}) = \pi^* \Omega$ where $i : P_{\vec{j}}^{(0)} \rightarrow P_{\vec{j}}$ is the inclusion map and $\pi : P_{\vec{j}}^{(0)} \rightarrow P_{\vec{j}}^{(0)} / \text{SU}(2)$ is the quotient map.

⁵ Explicitly this gives

$$N(z) = \left(\frac{1-|z|^2}{1+|z|^2} \tau_3 - \frac{z}{1+|z|^2} \tau_+ - \frac{\bar{z}}{1+|z|^2} \tau_- \right), \quad \tau_{\pm} = (\tau_1 \pm i\tau_2). \quad (11)$$

The τ algebra is

$$[\tau_1, \tau_2] = \tau_3 \quad \text{or} \quad [\tau_+, \tau_-] = \frac{2\tau_3}{i}, \quad [\tau_3, \tau_{\pm}] = \frac{\pm \tau_{\pm}}{i}.$$

⁶ Observe that $i_X(dz \wedge d\bar{z}) = X^z d\bar{z} - X^{\bar{z}} dz$.

these coordinates:

$$h_3(z) = j \frac{1 - |z|^2}{1 + |z|^2}, \quad \widehat{X}_3 = i(z\partial_z - \bar{z}\partial_{\bar{z}}), \quad (14)$$

$$h_+(z) = j \frac{-2\bar{z}}{1 + |z|^2}, \quad \widehat{X}_+ = i(\partial_z + \bar{z}^2\partial_{\bar{z}}), \quad (15)$$

$$h_-(z) = j \frac{-2z}{1 + |z|^2}, \quad \widehat{X}_- = -i(z^2\partial_z + \partial_{\bar{z}}). \quad (16)$$

The existence of a complex structure J implies that we can canonically extend the $SU(2)$ action to an action of $SL(2, \mathbb{C})$ on S^2 . That is, we have a mapping between the complexification of $\mathfrak{su}(2)$ and the space of *real* vector fields on S^2 given by $Y_1 + iY_2 \mapsto \widehat{Y}_1 + J(\widehat{Y}_2)$. It can be verified that this map is a Lie algebra morphism. The complex structure determines also a positive definite metric

$$g_j(\widehat{Y}_1, \widehat{Y}_2) \equiv \omega_j(J(\widehat{Y}_1), \widehat{Y}_2). \quad (17)$$

Explicitely this metric is

$$g_j(\widehat{Y}_1, \widehat{Y}_2) = j \{(Y_1|Y_2) - (Y_1|N)(N|Y_2)\}. \quad (18)$$

We can integrate the infinitesimal $SU(2)$ action (14,15,16) to a group action $e^{\widehat{Y}} \cdot f(N) = f(e^{-Y}Ne^Y)$ on functions. Defining $N(z^g) \equiv gN(z)g^{-1}$ one obtains that

$$z^g \equiv \frac{az + b}{\bar{a} - \bar{b}z} \quad \text{with} \quad g = \begin{pmatrix} a & b \\ -\bar{b} & \bar{a} \end{pmatrix}. \quad (19)$$

This is most naturally seen using the correspondence between the adjoint action on $N(z)$ and the left multiplication on $n(z)$:

$$gn(z) = n(z^g) \begin{pmatrix} \frac{a-b\bar{z}}{|a-b\bar{z}|} & 0 \\ 0 & \frac{\bar{a}-\bar{b}z}{|a-b\bar{z}|} \end{pmatrix}. \quad (20)$$

Thus, the $SL(2, \mathbb{C})$ action on S^2 is given by fractional transformations

$$N(z) \mapsto N(z^g), \quad \text{where} \quad z^g = \frac{az + b}{cz + d} \quad \text{and} \quad g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}. \quad (21)$$

The $SL(2, \mathbb{C})$ action on functions is $g \cdot f(z) = f(z^{g^{-1}})$.

We can now come back to the space of 4-gons and state the first main result that follows from the deep and beautiful work of Guillemin and Sternberg [19]. We have just shown that one has a natural action of $SL(2, \mathbb{C})$ on $P_{\vec{j}}$. As we have seen, the $SU(2)$ subgroup preserves the constrained set $P_{\vec{j}}^{(0)}$. On the contrary, the flow generated by the “imaginary” elements $J(\widehat{Y})$ is transverse to it. This follows from

$$J(\widehat{Y})h_Y(X) = -\omega_j(\widehat{Y}, J(\widehat{Y})) = g_j(\widehat{Y}, \widehat{Y}) > 0. \quad (22)$$

This suggests that we can relax the constraint as long as we ask for invariance under the complexified group $\mathrm{SL}(2, \mathbb{C})$ and this is the essence of the classical part of Guillemin and Sternberg's theorem. Namely, if one defines

$$P_{\vec{j}}^s \equiv \{(X(z_1), \dots, X(z_4)) \in P_{\vec{j}}^{(0)} \mid z_i \neq z_j \forall i \neq j\}, \quad (23)$$

this is a subset of the unconstrained phase space called the “stable” set. The complement of this set is the set of degenerate polygons and it is of measure 0.

Then, we have the non-trivial isomorphism [19, 27]

$$I : \mathcal{P}_{\vec{j}} = P_{\vec{j}}^{(0)} / \mathrm{SU}(2) \rightarrow P_{\vec{j}}^s / \mathrm{SL}(2, \mathbb{C}). \quad (24)$$

This isomorphism is quite surprising at first sight: it implies that we can identify a constrained phase space with an *unconstrained* one provided we demand invariance under the complexification of the gauge group.

First, one can show that we have an isomorphism

$$P_{\vec{j}}^s / \mathrm{SL}(2, \mathbb{C}) \rightarrow \mathbb{C} \setminus \{0, 1\}, \quad (25)$$

$$[z_1, \dots, z_4] \mapsto Z \equiv \frac{z_{41} z_{23}}{z_{43} z_{21}}, \quad (26)$$

where we have denoted $z_{ij} \equiv z_i - z_j$, the bracket stands for the equivalence class under $\mathrm{SL}(2, \mathbb{C})$ and Z is the cross-ratio. This is clear, since we can construct the fractional transformation

$$z \mapsto z^g = \frac{(z - z_1) z_{23}}{(z - z_3) z_{21}},$$

which is invertible and maps $(z_1, \dots, z_4)$ to $(0, 1, \infty, Z)$, and no fractional transformation except the identity maps $(0, 1, \infty, Z)$ to itself. The condition $z_i \neq z_j$ implies that $Z \neq 0, 1, \infty$.

The map I is then given by $I(z_1, \dots, z_4) = (0, 1, \infty, Z)$. The proof that this map is surjective, and hence an isomorphism, is non-trivial. It follows mainly from the work of Guillemin and Sternberg, but also from the specific study of invariant sections. The proof will be given in section V, since it requires more results.

B. The space of shape of tetrahedra

In this section we recall, following the seminal work of Barbieri [13], that the Polygon space $\mathcal{P}_{\vec{j}}$ is a suitable completion of the space of tetrahedra whose face areas are fixed to be equal to j_i .

One first needs to start with a subset of $H^{-1}(0)$ consisting of “nondegenerate” configurations, where a degenerate configuration of $H^{-1}(0)$ is such that $(X_1, \dots, X_4)$ is of rank at most 2, or equivalently, such that $\det(X_1, X_2, X_3) = 0$. If a configuration is degenerate, this means that all the vectors $X_i = j_i N(z_i)$ are orthogonal to a fixed direction. Since we are only interested in the polygon space, or the space $P_{\vec{j}}^{(0)}$ modulo $\mathrm{SU}(2)$, we can always choose this direction to be represented by τ_3 . The condition $\mathrm{tr}(\tau_3 N(z)) = 0$ is the condition that the z_i are real, hence the cross ratio Z labelling elements of $H^{-1}(0)/\mathrm{SU}(2)$ is real. Therefore, any nondegenerate configuration, which also corresponds to a nondegenerate tetrahedron,

as we are about to see, is labelled by a non-real cross-ratio. Thus, if we call the space of nondegenerate tetrahedra modulo rotations $\text{Tet}_{\vec{j}}$, we have

$$\text{Tet}_{\vec{j}} \simeq \mathbb{C} \setminus \mathbb{R}. \quad (27)$$

The fact that this space contains two disconnected components follows from the fact that, given a set of edge lengths, there are two possible tetrahedra that can be constructed and differ by a choice of orientation. Real cross ratios can be obtained as a limit of purely complex cross ratios. Such limits correspond to singular tetrahedra with zero volume and this shows that $\mathcal{P}_{\vec{j}}$ is a completion of the space of tetrahedra that includes all types of singular or degenerate ones:

$$\mathcal{P}_{\vec{j}} = \overline{\text{Tet}_{\vec{j}}} \quad (28)$$

If $\vec{j}$ is singular, i.e. $j_1 \pm j_2 \pm j_3 \pm j_4 \neq 0$ for some choice of sign, the closure of nondegenerate configurations in $\text{Tet}_{\vec{j}}$ could also contain cross ratios with values 0, 1 or ∞ .

In order to show this correspondence let us start from a nondegenerate configuration $(X_1, \dots, X_4) \in H^{-1}(0)$ and let us define

$$\ell^i \equiv \frac{1}{4V(X)} \epsilon^{ijkl} u_j [X_k, X_l], \quad V(X) = -2\text{tr}(X_1[X_2, X_3]), \quad (29)$$

where $V(X)$ is proportional to the square volume of the tetrahedron⁷, $u_i \equiv \frac{1}{2}(1, 1, 1, 1)$ is an auxiliary unit 4-vector whose components are all equal and ϵ is the totally antisymmetric tensor with four indices. These vectors satisfy a closure condition and a duality relation with X :

$$\sum_{i=1}^4 \ell^i = 0, \quad (\ell^i | X_j) = \delta_j^i - u^i u_j. \quad (30)$$

We can recover X_i from these vectors by a dual relation

$$X_i = \frac{1}{4V(\ell)} \epsilon_{ijkl} u^j [\ell^k, \ell^l], \quad V(X)V(\ell) = \frac{1}{4}, \quad (31)$$

with $V(\ell) = \det(\ell_1, \ell_2, \ell_3)$. We can therefore construct a tetrahedron whose area normal vectors are exactly given by the X_i . The edge vectors of this tetrahedron are

$$L^{ij} \equiv \sqrt{|V(X)|} (\ell^i - \ell^j). \quad (32)$$

That is, X_i is the vector normal to the face opposite to the vertex i and its norm is twice the area of this face⁸. The relationship between the edge vectors L^{ij} and the X_i is

$$X_1 = \epsilon [L^{24}, L^{34}] \text{ etc. }, \quad \text{where } V(L) \equiv \det(L^{14}, L^{24}, L^{34}) = \epsilon \sqrt{|V(X)|}, \quad (33)$$

and $\epsilon = \text{sgn } V(X)$ characterises the orientation of the tetrahedron. The proof of these statements can be obtained by direct computation and a 4-dimensional analogue is given in

⁷ If one uses the $\mathbb{R}^3$ notation for X instead of $\mathfrak{su}(2)$, we have $V = \det(X_1, X_2, X_3)$.

⁸ A priori, there are two tetrahedra of opposite orientation that have the same area normal vectors X_i . With definition (32) we choose one of these two tetrahedra: namely, the one for which $\text{sgn } \det(L^{14}, L^{24}, L^{34}) = \text{sgn } V(X)$.

the appendix of [5]. The relations between the volumes shows that when X_i is a sequence of nondegenerate configurations converging toward a degenerate one, then $V(X) \rightarrow 0$ thus the volume of the tetrahedra $V(L)$ also goes to 0. Note that at the same time $V(\ell) \rightarrow \infty$.

To complete this geometrical study it is interesting to translate the canonical variables and the cross ratio into the tetrahedral language. In order to do so we denote by θ_{ij} the dihedral angle of the edge shared by the face i and j , i.e. $\cos \theta_{ij} = (N_i | N_j)$. The first canonical variable j_{12} is related to the dihedral angle θ_{12} by

$$j_{12}^2 = j_1^2 + j_2^2 + 2 \cos \theta_{12}, \quad (34)$$

while the “bending” angle ϕ_{12} is the angle between opposite edges (12) and (34), i.e.

$$\cos \phi_{12} = \frac{(\ell^{12} | \ell^{34})}{|\ell^{12}| |\ell^{34}|}. \quad (35)$$

More generally, let us denote by ϕ_{jk}^i the angle between the edges (ij) and (ik) , that is, $\cos \phi_{jk}^i = (\ell^{ij} | \ell^{ik}) / |\ell^{ij}| |\ell^{ik}|$. We can express the cross ratio, which parametrises all the possible tetrahedral shapes, in terms of these geometrical data:

$$Z = \frac{\sin \frac{\theta_{41}}{2} \sin \frac{\theta_{23}}{2}}{\sin \frac{\theta_{43}}{2} \sin \frac{\theta_{21}}{2}} e^{i(\phi_{13}^4 + \phi_{13}^2 - \phi_{24}^1 - \phi_{24}^3)}. \quad (36)$$

III. THE QUANTUM GEOMETRY OF COHERENT STATES

In this section we recall some definitions of coherent states and their relations with geometric quantisation. We refer the reader to the textbook [30] for more information on these states.

The Lie algebra generators of $SU(2)$ are denoted J_i and satisfy the algebra

$$[J_+, J_-] = 2J_3, \quad [J_3, J_\pm] = \pm J_\pm, \quad (37)$$

and the reality conditions $J_+^\dagger = J_-$, $J_3^\dagger = J_3$. The spin j representation of this algebra is

$$\begin{aligned} T^j(J_+) |j, m\rangle &= \sqrt{(j+m+1)(j-m)} |j, m+1\rangle, \\ T^j(J_3) |j, m\rangle &= m |j, m\rangle, \\ T^j(J_-) |j, m\rangle &= \sqrt{(j-m+1)(j+m)} |j, m-1\rangle, \end{aligned} \quad (38)$$

with Casimir $J^2 = j(j+1)$ and we denote by V^j the corresponding carrier space. Note that in the spin $1/2$ representation we have $J_a = \sigma_a/2 = i\tau_a$. Given such a representation, we define the group element $n(z)$ as in eq. (12). In terms of Lie algebra elements $n(z)$ can be expressed as

$$n(z) = e^{zJ_+ - \bar{z}J_-}. \quad (39)$$

The group coherent states are defined by

$$|j, n(z)\rangle \equiv T^j(n(z)) |j, -j\rangle. \quad (40)$$

In the following section it will be convenient to label coherent states not only in terms of group elements n , but also directly by the vector N in S^2 which n represents. That is, we define a coherent state labelled by N :

$$|j, N\rangle \equiv |j, n(N)\rangle, \quad \text{with} \quad n(N) = \frac{1}{\sqrt{2(1+N_3)}} \begin{pmatrix} 1+N_3 & -N_1-iN_2 \\ N_1+iN_2 & 1+N_3 \end{pmatrix} \quad (41)$$

In this way, $|j, N(z)\rangle = |j, n(z)\rangle$.

One key property of these states is their coherence under the tensor product, that is,

$$|j, n(z)\rangle = |n(z)\rangle^{\otimes 2j}, \quad \text{where} \quad |n(z)\rangle \equiv |1/2, n(z)\rangle. \quad (42)$$

To emphasize this coherence property we will preferably use this notation for the coherent state. The second key property is the property of holomorphicity:

$$|n(z)\rangle^{\otimes 2j} = \frac{1}{(1+|z|^2)^j} |z\rangle^{\otimes 2j}, \quad \text{where} \quad |z\rangle \equiv e^{zJ_+} |1/2, -1/2\rangle. \quad (43)$$

From this one can compute the hermitian product of such states,

$$^{\otimes 2j} \langle n(w) | n(z) \rangle^{\otimes 2j} = \langle n(w) | n(z) \rangle^{2j} = \frac{(1+\bar{w}z)^{2j}}{(1+|w|^2)^j (1+|z|^2)^j}, \quad (44)$$

and check the completeness relation

$$\mathbb{1}_j = \frac{d_j}{2\pi} \int \frac{d^2 z}{(1+|z|^2)^2} (|n(z)\rangle \langle n(z)|)^{\otimes 2j} = d_j \int_{S^2} d^2 n(z) (|n(z)\rangle \langle n(z)|)^{\otimes 2j}, \quad (45)$$

where $d_j = 2j+1$ and $d^2 n(z)$ is the normalised measure on the unit sphere. From the definition of the coherent state and the behavior of $n(z)$ under left multiplication expressed in (20)⁹ we can get the expression of the SU(2) group action on the group coherent state:

$$T^j(g) |n(z)\rangle^{\otimes 2j} = |gn(z)\rangle^{\otimes 2j} = \left(\frac{\bar{a} - \bar{b}z}{|\bar{a} - \bar{b}z|} \right)^{2j} |n(z^g)\rangle^{\otimes 2j}, \quad z^g \equiv \frac{az+b}{-\bar{b}z+\bar{a}} \quad \text{if} \quad g = \begin{pmatrix} a & b \\ -\bar{b} & \bar{a} \end{pmatrix}.$$

Using that $|\bar{a} - \bar{b}z|^2(1+|z^g|^2) = (1+|z|^2)$ if $g \in \text{SU}(2)$, we can get the corresponding action for the holomorphic state $|z\rangle$ which reads

$$T^j(g) |z\rangle^{\otimes 2j} = (\bar{a} - \bar{b}z)^{2j} |z^g\rangle^{\otimes 2j}. \quad (46)$$

The essential point is that this action is purely holomorphic, since $\partial_{\bar{z}} |z\rangle^{\otimes 2j} = 0$. Therefore, as in the classical setting, we can extend the SU(2) action to an action of SL(2, $\mathbb{C}$). That is,

$$T^j(g) |z\rangle^{\otimes 2j} = (cz+d)^{2j} |z^g\rangle^{\otimes 2j}, \quad z^g \equiv \frac{az+b}{cz+d} \quad \text{if} \quad g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}(2, \mathbb{C}). \quad (47)$$

⁹ Note that we can write this relation as $gn(z) = n(z^g) \left(\frac{\bar{a}-\bar{b}z}{|\bar{a}-\bar{b}z|} \right)^{-2J_3}$.

We can lift this action back from the holomorphic to the group coherent state and finally get

$$\begin{aligned} T^j(g)|n(z)\rangle^{\otimes 2j} &= \left(\frac{cz+d}{|cz+d|}\right)^{2j} \left(\frac{|cz+d|^2(1+|z^g|^2)}{1+|z|^2}\right)^j |n(z^g)\rangle^{\otimes 2j} \\ &= \left(\frac{cz+d}{|cz+d|}\right)^{2j} \left(\frac{\langle z|g^\dagger g|z\rangle}{\langle z|z\rangle}\right)^j |n(z^g)\rangle^{\otimes 2j} \\ &\equiv \rho_g^{-j}(z)|n(z^g)\rangle^{\otimes 2j}. \end{aligned} \quad (48)$$

$$\equiv \rho_g^{-j}(z)|n(z^g)\rangle^{\otimes 2j}. \quad (49)$$

In the second line we used g interchangeably for $T^{1/2}(g)$. We have introduced a pairing $\rho_g(z)$ between an element of $\text{SL}(2, \mathbb{C})$ and an element of S^2 which is explicitly given by

$$\rho_g(z) \equiv \frac{1+|z|^2}{|cz+d|^2+|az+b|^2} \left(\frac{|cz+d|}{cz+d}\right)^2, \quad \text{for } g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}(2, \mathbb{C}). \quad (50)$$

From (47) one can derive explicitly the action of infinitesimal $\text{SU}(2)$ generators acting on the holomorphic states:

$$\begin{aligned} \langle s|T^j(J_+)|z\rangle^{\otimes 2j} &= \partial_z \langle s|z\rangle^{\otimes 2j}, \\ \langle s|T^j(J_3)|z\rangle^{\otimes 2j} &= (z\partial_z - j) \langle s|z\rangle^{\otimes 2j}, \\ \langle s|T^j(J_-)|z\rangle^{\otimes 2j} &= (-z^2\partial_z + 2jz) \langle s|z\rangle^{\otimes 2j}. \end{aligned} \quad (51)$$

Here, $|s\rangle$ is any state in the spin j representation. From this we can now easily compute the expectation value of Lie algebra generators and find that it is simply given by the moment map h_a :

$$\frac{\langle j, z|T^j(J_a)|j, z\rangle}{\langle j, z|j, z\rangle} = -jN_a(z) = -h_a(z) \quad (52)$$

We will also need the expression for the action of a general vector field $\widehat{X}$ on the moment map: for $X \in \mathfrak{sl}(2, \mathbb{C})$ and $Y \in \mathfrak{su}(2)$

$$e^{-\widehat{X}} \cdot h_Y(z) = \frac{1}{i} \frac{\langle j, z|T^j(e^{X^\dagger} Y e^X)|j, z\rangle}{\langle j, z|T^j(e^{X^\dagger} e^X)|j, z\rangle}. \quad (53)$$

This follows directly from the fact that on the one hand $e^{-\widehat{X}} \cdot h_Y(z) = h_Y(z^g)$ with $g = e^X$, and on the other hand $h_Y(z^g)$ can be expressed as the right-hand side of (53) using the coherent state transformation (47) and the relation (52) between the Hamiltonian and the expectation value (recall that $J_a = i\tau_a$).

We can also see that in the case $X \in \mathfrak{su}(2)$ (that is, X anti-hermitian) the previous expression reduces to the action (5) of $\text{SU}(2)$ on functions on S^2 . By differentiating the previous expression we also get a relation between the metric and the symmetric connected 2-point function: given $X, Y \in \mathfrak{su}(2)$ we have

$$\begin{aligned} -g_j(\widehat{X}, \widehat{Y}) &= \frac{\langle j, z|T^j(XY + YX)|j, z\rangle}{\langle j, z|j, z\rangle} - 2 \frac{\langle j, z|T^j(X)|j, z\rangle}{\langle j, z|j, z\rangle} \frac{\langle j, z|T^j(Y)|j, z\rangle}{\langle j, z|j, z\rangle} \\ &= -jX^a(\delta_{ab} - N_a(z)N_b(z))Y^b \end{aligned} \quad (54)$$

with $X = X^a \tau_a$. More generally, the connected two point function is related to the hermitian form

$$-g_j(\widehat{X}, \widehat{Y}) + i\omega_j(\widehat{X}, \widehat{Y}) = 2 \frac{\langle j, z | T^j(XY) | j, z \rangle}{\langle j, z | j, z \rangle} - 2 \frac{\langle j, z | T^j(X) | j, z \rangle}{\langle j, z | j, z \rangle} \frac{\langle j, z | T^j(Y) | j, z \rangle}{\langle j, z | j, z \rangle} \quad (55)$$

$$= -jX^a (\delta_{ab} - N_a(z)N_b(z) - i\epsilon_{abc}N^c(z)) Y^b. \quad (56)$$

A. Geometric quantisation

The previous description of coherent states is the usual one used in the physical literature. In the mathematical literature [20, 22, 23] one uses preferably the language of geometric quantisation to describe the same construction. For the reader's convenience and in order to connect the mathematical and physical terminology we describe the relationship between these two languages in a pedestrian manner.

Firstly, in order to define the geometrical quantisation we need a Hermitian line bundle L over S^2 of curvature $i\omega_j$. A natural hermitian line bundle over S^2 is given by $\mathbb{CP}_1 \rightarrow S^2$. Denoting this line bundle by L , all the other line bundles are obtained by tensorisation $L_j \equiv L^{\otimes 2j}$. The conditions on this line bundle mean that we have a covariant derivative $\nabla = d + A$, with hermitian connection and curvature given by

$$A_z = -j \frac{\bar{z}}{1 + |z|^2}, \quad A_{\bar{z}} = j \frac{z}{1 + |z|^2}, \quad (57)$$

$$F_{z\bar{z}} = \partial_z A_{\bar{z}} - \partial_{\bar{z}} A_z = \frac{2j}{(1 + |z|^2)^2}. \quad (58)$$

In geometric quantisation one identifies the representation space of spin j with the space of holomorphic sections of L_j denoted $H^0(S^2, L_j)$. Explicitly, the space of sections of L_j is the space of functions $s(z, \bar{z})$ such that both s and $\tilde{s}(z) = \frac{z^{2j}}{|z|^{2j}} s(-\frac{1}{z})$ are holomorphic sections. That is, they are analytic and solutions of $\nabla_{\widehat{X} - iJ(\widehat{X})} s = 0$. This implies that

$$s(z, \bar{z}) = \frac{\hat{s}(z)}{(1 + |z|^2)^j},$$

where $\hat{s}(z)$ is a holomorphic polynomial of degree at most $2j$.

$H^0(S^2, L_j)$ carries a representation of $SU(2)$ given by the differential operators

$$Y \cdot s = (\nabla_{\widehat{Y}} - ih_Y) s \quad (59)$$

where Y is an anti-hermitian element and h_Y is the momentum map associated with the $SU(2)$ action on S^2_j . The fact that this forms a representation follows directly from the symplectic identities

$$\omega(\widehat{X}, \widehat{Y}) = -iF(\widehat{X}, \widehat{Y}) = \widehat{X}h_Y = h_{[X, Y]}. \quad (60)$$

The link between this representation and the previous coherent state representation can be described as follows. Suppose that $|s\rangle$ is a state in V_j and define

$$s(z) \equiv \langle s | n(z) \rangle^{\otimes 2j}. \quad (61)$$

One can verify that this is a holomorphic section of L_j . In particular, $\nabla_{\bar{z}}s(z) = 0$, which follows from

$$\nabla_{\bar{z}}s(z) = \frac{\partial_{\bar{z}}\hat{s}(z)}{(1+|z|^2)^j} = 0, \quad \hat{s}(z) \equiv \langle s|j, z \rangle. \quad (62)$$

Furthermore, the geometric representations coincide with the group coherent state representation:

$$Y \cdot s(z) = (\nabla_{\widehat{Y}} - i h_Y) s(z) = -\langle s|T^j(Y)|n(z)\rangle^{\otimes 2j} = -\frac{\langle s|T^j(Y)|z\rangle^{\otimes 2j}}{\langle z|z\rangle^{\otimes 2j}} \quad (63)$$

This follows from the fact that the “Lagrangians” $L_X \equiv \theta(\widehat{X}) - h_X$ associated with the holomorphic symplectic potential $\theta \equiv 2ij \frac{\bar{z}dz}{1+|z|^2}$ coincide with the scalar terms in (51), that is,

$$L_{X_3} = -j, \quad L_{X_+} = 0, \quad L_{X_-} = 2jz. \quad (64)$$

The language of geometric quantisation explains also the extension of the $SU(2)$ action to $SL(2, \mathbb{C})$. Firstly, the complex structure allows us to extend the mapping $X \rightarrow \widehat{X}$ from Lie algebra to real vector fields to hermitian operators by $(i\widehat{X}) = J(\widehat{X})$. Then, the holomorphicity of the section implies that $\nabla_{J(\widehat{X})} = i\nabla_{\widehat{X}}$, hence

$$(iY) \cdot s(z) = \left(\nabla_{(iY)} - i h_{iY} \right) s(z) = \left(\nabla_{J(\widehat{Y})} + h_Y \right) s(z). \quad (65)$$

IV. QUANTISATION COMMUTES WITH REDUCTION

Now that we have described all the necessary ingredients we can focus on our main task: the construction of the quantisation of the space of shape of tetrahedra, which is the 4-gon space $\mathcal{P}_{\vec{j}}$. One route to the quantisation of this space is the route proposed by Barbieri [13] in his seminal work and studied since then in the quantum gravity literature: according to this scheme one first quantises the phase space of four vectors $X_i \in S_{j_i}^2$, promotes these vectors to operators associated to 4 copies of $\mathfrak{su}(2)$, where in each case $X_i^2 = j_i(j_i + 1)$, and *then* one imposes the constraints $\sum_i X_i = 0$ at the operatorial level. The Hilbert space for this quantisation is the space of 4-valent intertwiners

$$H_{\vec{j}} = (V^{j_1} \dots \otimes V^{j_4})^{SU(2)}, \quad (66)$$

i.e. the space of $SU(2)$ invariant vectors. A basis for this space is given by invariant states $|j_i, i_{12}\rangle$ diagonalising the operators $(X_1 + X_2)^2$ with eigenvalue $i_{12}(i_{12} + 1)$ and labelled by an intertwiner label i_{12} which is a spin. The completeness relation for these states is $\mathbb{1}_{\vec{j}} = \sum_i d_i |j_i, i\rangle \langle j_i, i|$. Recently, it has been advocated by Livine and Speziale [17] that a more convenient basis is the “coherent intertwiner” basis. For this basis one starts with a product of 4 coherent states and then averages them with respect to the group action in order to get an invariant state:

$$||\vec{j}, N_i\rangle \equiv \int_{SU(2)} dk \otimes_{j_i} T^{j_i}(k) |j_i, N_i\rangle \quad (67)$$

In this basis the resolution of identity reads

$$\mathbb{1}_{\vec{j}} = \prod_i d_{j_i} \int \prod_i d^2 N_i ||\vec{j}, N_i\rangle \langle \vec{j}, N_i||,$$

where $\mathbb{1}_{\vec{j}}$ denotes the identity in (or projector onto) $H_{\vec{j}}$ and d^2N is the normalised measure on S^2 .

Another route toward the quantisation of the same space, which is conceptually much simpler, is to first impose the constraints and then quantise. As we have seen, the reduced space $\mathcal{P}_{\vec{j}} = P_{\vec{j}}^{(0)}/\text{SU}(2)$ is a symplectic manifold and we even know some nice symplectic coordinates (j_{12}, ϕ_{12}) on this space (see (9)). All $\text{SU}(2)$ invariant function can be expressed as a function of these variables and the quantisation in this real polarisation is straightforward: the Hilbert space $H_{\vec{j}}^{(0)}$ is the space of L^2 function of ϕ_{12} (say) and the completeness relation reads $\mathbb{1}_{\vec{j}} = \int_0^{2\pi} d\phi_{12} |\phi_{12}\rangle \langle \phi_{12}|$.

We have also seen that $\mathcal{P}_{\vec{j}}$, being the quotient of a Kähler manifold¹⁰, is itself a Kähler manifold. The complex coordinate associated with this Kähler structure is the cross ratio $Z(z_i)$ (see eq. (26)). Following the procedure of geometric quantisation presented in the previous section, we can construct the coherent states $|Z\rangle$ associated to this complex coordinate. The completeness relation for these states is

$$\mathbb{1}_{\vec{j}} = \int_{\mathcal{P}_{\vec{j}}} \Omega(Z) |Z\rangle \langle Z|$$

with Ω being the symplectic structure on $\mathcal{P}_{\vec{j}}$.

The key and essential result following from the general framework of Guillemin–Sternberg [19] and the detailed study of its unitarity properties by B. Hall [38] is that one should expect an isomorphism between the two Hilbert spaces $\hat{I} : H_{\vec{j}}^{(0)} \rightarrow H_{\vec{j}}$. This isomorphism is the mathematical translation of the statement that “Quantisation commutes with reduction”. The purpose of section V is to give a proof of this isomorphism *and* to construct it explicitly. As we will see, the correspondence is such that the “reduced” coherent state $|Z\rangle$ maps, up to an overall normalisation, to a group-averaged coherent state, labelled by a point on the constraint surface:

$$I(|Z\rangle) = \tilde{\rho}_{\vec{j}}(z_i) ||\vec{j}, N_i\rangle \quad \text{for all } z_i \text{ s.t. } Z = Z(z_i) \text{ and } \sum_i j_i N(z_i) = 0. \quad (68)$$

$\tilde{\rho}_{\vec{j}}(z_i)$ is a strictly positive function invariant under the action of $\text{SL}(2, \mathbb{C})$. The proof we are going to give is self-contained and does not require any prior knowledge of geometric quantisation and the work of Guillemin–Sternberg.

A. 3- and 4-valent section

Before going into the proof of our main statement we first describe in more detail the space of 3- and 4-valent invariant sections and some properties of the $\text{SU}(2)$ invariant group coherent state (67). Starting from this state it will be convenient to define the corresponding holomorphic state

$$||\vec{j}, z_i\rangle \equiv \prod_i (1 + |z_i|^2)^{j_i} ||\vec{j}, N(z_i)\rangle. \quad (69)$$

¹⁰ Recall that a Kähler manifold is a symplectic manifold with a complex structure compatible with the symplectic form, i.e. $\omega(\hat{X}, \hat{Y}) = \omega(J(\hat{X}), J(\hat{Y}))$, and $B(\hat{X}, \hat{Y}) = \omega(J(\hat{X}), \hat{Y})$ is a positive definite pairing.

Given a state $|S\rangle$ in $H_j = (V^{j_1} \otimes \dots \otimes V^{j_4})^{\text{SU}(2)}$, we can construct the “invariant holomorphic section”

$$S^{\vec{j}}(z_i) \equiv \langle S | |\vec{j}, z_i \rangle. \quad (70)$$

Such sections are characterised by the fact that they are polynomials of degree $2j_i$ in the variable z_i and by their property of invariance under $\text{SL}(2, \mathbb{C})$ transformations: namely,

$$S^{\vec{j}}(z_i + a) = S^{\vec{j}}(z_i), \quad \prod_i z_i^{2j_i} S^{\vec{j}}\left(-\frac{1}{z_i}\right) = S^{\vec{j}}(z_i), \quad S^{\vec{j}}(\lambda Z_i) = \lambda^{2j_i} S^{\vec{j}}(z_i). \quad (71)$$

Conversely, any such section defines a state in $H_j = (V^{j_1} \dots \otimes V^{j_4})^{\text{SU}(2)}$ by

$$|S\rangle = \prod_i d_{j_i} \int \prod_i d^2 N(z_i) |\vec{j}, N(z_i)\rangle \frac{\overline{S^{\vec{j}}(z_i)}}{(1 + |z_i|^2)^{j_i}}. \quad (72)$$

Let us first look at the space of invariant sections which depends only on three entries (z_1, z_2, z_3) . The conditions of $\text{SL}(2, \mathbb{C})$ invariance (71) fix the form of the section uniquely up to a normalisation. This unique section is given by

$$S^{j_1 j_2 j_3}(z_1, z_2, z_3) = \sqrt{N_{j_1 j_2 j_3}} (z_1 - z_2)^{\Delta_3} (z_2 - z_3)^{\Delta_1} (z_3 - z_1)^{\Delta_2}, \quad (73)$$

where $\Delta_i = j_1 + j_2 + j_3 - 2j_i$ and N_{j_i} is a normalisation factor chosen such that

$$1 = \langle S | S \rangle = \prod_{i=1}^3 d_{j_i} \int \prod_{i=1}^3 d^2 N(z_i) \frac{|S^{j_1 j_2 j_3}(z_1, z_2, z_3)|^2}{\prod_{i=1}^3 (1 + |z_i|^2)^{2j_i}}. \quad (74)$$

It is explicitly given by (see [31])

$$N_{j_1 j_2 j_3}^{-1} = \frac{[j_1 + j_2 + j_3 + 1]! [-j_1 + j_2 + j_3]! [j_1 - j_2 + j_3]! [j_1 + j_2 - j_3]!}{[2j_1]! [2j_2]! [2j_3]!}. \quad (75)$$

Next we look at the space of invariant sections which depend on four entries (z_1, z_2, z_3, z_4) . The condition of $\text{SL}(2, \mathbb{C})$ invariance no longer fixes the functional form of these sections uniquely. It implies, however, that they have the form

$$S(z_i) = z_{12}^{j_1 + j_2 - j_3 + j_4} z_{23}^{(-j_1 + j_2 + j_3 - j_4)} z_{31}^{(j_1 - j_2 + j_3 - j_4)} z_{34}^{2j_4} s(Z), \quad (76)$$

where Z is the cross ratio

$$Z \equiv \frac{z_{41} z_{23}}{z_{43} z_{21}}, \quad z_{ij} \equiv z_i - z_j, \quad (77)$$

and $s(Z) = \lim_{X \rightarrow \infty} X^{-2j_3} S(0, 1, X, Z)$. This section should also be a polynomial in z_i of degree at most $2j_i$. Therefore, s should be a polynomial in Z of degree smaller than $\min(2j_4, j_4 + j_1 + j_2 - j_3)$ and of valuation at least $\max(0, j_4 - j_1 + j_2 - j_3, j_4 + j_1 - j_2 - j_3)$.

The usual spin network basis $|\vec{j}, i\rangle$ corresponds to a section denoted $S_i^{\vec{j}}(z_i) \equiv \langle \vec{j}, i | |\vec{j}, z_i \rangle$. This section diagonalises the operator $\Delta_{12} \equiv X^{(1)} \cdot X^{(2)}$ with eigenvalue $i(i+1) - j_1(j_1+1) - j_2(j_2+1)$, where $X^{(i)}$ denotes the action on the variable z_i . This differential operator is given by

$$\Delta_{12} = -\frac{1}{2} z_{12}^2 \partial_1 \partial_2 + z_{12} (j_1 \partial_2 - j_2 \partial_1) + j_1 j_2. \quad (78)$$

The projection of the coherent state on the spin network basis can also be expressed as an integral involving the product of two trivalent invariant (Clebsch–Gordan) sections:

$$S_i^{\vec{j}}(z_i) = d_i \int \frac{S^{j_1 j_2 i}(z_1, z_2, z) S^{i j_3 j_4}(-\bar{z}^{-1}, z_3, z_4)}{(1 + |z|^2)^{2i}} z^{2i} d^2 N(z). \quad (79)$$

V. THE QUANTUM EQUIVALENCE

The purpose of this section is to first present some of the key properties satisfied by the coherent intertwiner, then to give a proof of the classical correspondence (24) and finally to state and prove our main result: the explicit isomorphism between the space of 4-valent intertwiners and the space generated by coherent states labelled by classical tetrahedra.

A. Some properties of invariant coherent states

As discussed previously, the coherent intertwiner (67) is defined by the projection of the usual group coherent states:

$$||\vec{j}, N_i\rangle \equiv \Pi^{(0)} \otimes_i |j_i, N_i\rangle, \quad \Pi^{(0)} \equiv \int_{\text{SU}(2)} dk \otimes_{j_i} T^{j_i}(k) \quad (80)$$

Here, $\Pi^{(0)}$ is a group averaging projector annihilating the diagonal $\text{SU}(2)$ action. Since the coherent state is holomorphic this invariance extends to the $\text{SL}(2, \mathbb{C})$ action. Together with the rules of transformation (49), the invariance property $\Pi^{(0)} (\otimes_i T^{j_i}(g)) = \Pi^{(0)}$ implies the following key transformation property: given any $g \in \text{SL}(2, \mathbb{C})$

$$||\vec{j}, N(z_i^g)\rangle = \rho_g^{\vec{j}}(z_i) ||\vec{j}, N(z_i)\rangle, \quad (81)$$

where the prefactor equals the product

$$\rho_g^{\vec{j}}(z_i) \equiv \prod_i \rho_g^{j_i}(z_i), \quad \rho_g(z) = \left(\frac{|cz + d|}{cz + d} \right)^2 \frac{\langle z|z \rangle}{\langle z|g^\dagger g|z \rangle}, \quad (82)$$

and $\langle z|z \rangle = 1 + |z|^2$. We also introduce a notation for the norm of the covariant intertwiner $||N(z_i)||_{\vec{j}}^2 \equiv \langle \vec{j}, N(z_i) | \vec{j}, N(z_i) \rangle$. The factor $\rho_{\vec{j}}$ is just the ratio of this norms

$$|\rho_g^{\vec{j}}|^2(z_i) = \frac{||N(z_i^g)||_{\vec{j}}^2}{||N(z_i)||_{\vec{j}}^2}. \quad (83)$$

Note that in the following we will also use the notation $\rho_g(N_i)$ with the obvious definition $\rho_g(N(z_i)) \equiv \rho_g(z_i)$. The prefactor $\rho_g^{\vec{j}}(z_i)$ controls the behavior of the invariant states along the $\text{SL}(2, \mathbb{C})$ orbits and satisfies a number of key properties that we now list:

Proposition V.1 *Suppose that (z_i) is a point in $P_{\vec{j}}$ with nonzero norm $||N(z_i)||_{\vec{j}}$.*

1- The first derivatives of $||N(z_i)||_{\vec{j}}$ for an anti-hermitian $X = X^a \tau_a$ are given by

$$\widehat{(iX)} \ln ||N(z_i)||_{\vec{j}} = -H_X(z_i), \quad (84)$$

$$\widehat{(iY)} \widehat{(iX)} \ln ||N(z_i)||_{\vec{j}} = -G(X, Y)(z_i), \quad (85)$$

where $H_X(N_i) = \sum_i j_i N_i$ is the Hamiltonian constraint and $G = \sum_i g_{j_i}$ is the metric on the orbits $\text{SL}(2, \mathbb{C}) \cdot (z_i)$. Explicitly,

$$G_{ab} \equiv G(\tau_a, \tau_b) = \sum_i j_i (\delta_{ab} - N_a(z_i) N_b(z_i)). \quad (86)$$

2- Suppose that $(N_i) \in P_{\vec{J}}^{(0)}$, i.e. $\sum_i j_i N_i = 0$, then $|\rho_{\vec{g}}^{\vec{J}}(N_i)| \leq 1$ for all $g \in \text{SL}(2, \mathbb{C})$. Moreover, the equality is satisfied iff $g \in \text{SU}(2)$.

3- We can relate the norm of $\rho_{\vec{g}}^{\vec{J}}(z_i)$ to the integral of the Hamiltonian constraint:

$$|\rho_{e^{iX}}^{\vec{J}}(N_i)|^2 = \exp \left(\int_0^1 dt e^{-t\widehat{X}} \cdot H_{(X-X^\dagger)}(z_i) \right), \quad (87)$$

with $X \in \mathfrak{sl}(2, \mathbb{C})$.

Let us start by proving point 3: given $X \in \mathfrak{sl}(2, \mathbb{C})$ we have

$$|\rho_{e^{iX}}^{\vec{J}}(z_i)|^2 = \prod_i \frac{\langle j_i, z_i | j_i, z_i \rangle}{\langle j_i, z_i | e^{-iX^\dagger} e^{iX} | j_i, z_i \rangle} \quad (88)$$

$$= \exp \left(- \int_0^1 dt \partial_t \sum_i \ln \langle j_i, z_i | e^{-itX^\dagger} e^{itX} | j_i, z_i \rangle \right) \quad (89)$$

$$= \exp \left(- \int_0^1 dt \sum_i \frac{\langle j_i, z_i | e^{-itX^\dagger} (-iX^\dagger + iX) e^{itX} | j_i, z_i \rangle}{\langle j_i, z_i | e^{-itX^\dagger} e^{itX} | j_i, z_i \rangle} \right) \quad (90)$$

$$= \exp \left(\int_0^1 dt H_{(X-X^\dagger)}(e^{itX} \cdot z_i) \right) = \exp \left(\int_0^1 dt e^{-tJ(\widehat{X})} \cdot H_{(X-X^\dagger)}(z_i) \right). \quad (91)$$

In the last equalities we have used the definition of H_X in terms of the expectation value given in section III. This expression coincides with the general expression derived by B. Hall [38]. Taking the logarithm of this formula and restricting to $X \in \mathfrak{su}(2)$ one obtains that

$$\ln(|\rho_{e^{iuX}}^{\vec{J}}(z_i)|) = \int_0^u dt e^{-tJ(\widehat{X})} \cdot H_X(z_i). \quad (92)$$

By taking the derivative of this expression with respect to u , one obtains that

$$\partial_u \ln(|\rho_{e^{iuX}}^{\vec{J}}(z_i)|) \Big|_{u=0} = H_X(z_i) = -J(\widehat{X}) \ln ||N(z_i)||_{\vec{J}}, \quad (93)$$

where the second equality follows from the relationship (83):

$$\ln(|\rho_{e^{iuX}}^{\vec{J}}(z_i)|) = \left(e^{-uJ(\widehat{X})} - 1 \right) \ln ||N(z_i)||_{\vec{J}} \quad (94)$$

By taking another derivative of (93) one obtains easily property 1, since $J(\widehat{X})H_Y = G(X, Y)$.

From the relation (81) one clearly sees that if $||N(z_i)||_{\vec{J}} \neq 0$, then for any $g \in \text{SL}(2, \mathbb{C})$ we also have $||N(z_i^g)||_{\vec{J}} \neq 0$, since $|\rho_{\vec{g}}^{\vec{J}}(z_i)|$ is always strictly positive. Let us consider such a point $(z_i) \in P_{\vec{J}}$. From property 1 we infer that if (z_i) is a critical point of $||N(z_i)||_{\vec{J}}$, then it necessarily lies on the constraint surface $P_{\vec{J}}^{(0)}$. Moreover, since the second derivative is always negative, this means that this critical point is a maximum. This proves property 2.

B. Back to the Guillemin–Sternberg isomorphism

We can in fact show more: consider a point $(z_i) \in P_{\vec{J}}$ with nonzero norm $||N(z_i)||_{\vec{J}} \neq 0$, and let us look at the closure of the orbit $O_{z_i} = \text{SL}(2, \mathbb{C}) \cdot (z_i)$. Since this is a closed

submanifold of a compact space and since we established that $\|N(z_i^g)\|_{\vec{j}}$ is a concave function of g , it reaches its maximum for a given element of the closure of O_{z_i} . Because of property 1 this maximum occurs in the constraint space $P_{\vec{j}}^{(0)}$. The orbit space $\text{SL}(2, \mathbb{C}) \cdot P_{\vec{j}}^{(0)}$ is an open neighborhood of $P_{\vec{j}}^{(0)}$, because at a given point $(z_i) \in P_{\vec{j}}^{(0)}$ the vectors $J(\hat{X})$ are all orthogonal to tangent vectors in $P_{\vec{j}}^{(0)}$. The latter follows from

$$G(\xi, J(\hat{X}))(z_i) = \xi H_X(z_i) = 0, \quad \text{when } (z_i) \in P_{\vec{j}}^{(0)}, \quad \xi \in T_{(z_i)} P_{\vec{j}}^{(0)}.$$

Since $\text{SL}(2, \mathbb{C}) \cdot P_{\vec{j}}^{(0)}$ is an open neighborhood of $P_{\vec{j}}^{(0)}$, the orbit O_{z_i} intersects it. Therefore, $z_i \in \text{SL}(2, \mathbb{C}) \cdot P_{\vec{j}}^{(0)}$.

This proves the fundamental result of Guillemin–Sternberg: namely, the characterisation of the orbit space

$$P_{\vec{j}}^s \equiv \text{SL}(2, \mathbb{C}) \cdot P_{\vec{j}}^{(0)} = \{z_i \in P_{\vec{j}} \mid \|N(z_i)\|_{\vec{j}} \neq 0\}$$

as the space, where the norm of the coherent intertwiner does not vanish¹¹.

We can now finally tighten one of our loose ends and get the final characterisation of the stable set $P_{\vec{j}}^s$. First we expand the coherent state norm in terms of the usual orthonormal intertwiner basis $|\vec{j}, i\rangle$ diagonalising the operator $(X_1 + X_2)$:

$$\langle \vec{j}, z | \vec{j}, z \rangle = \sum_i d_i |S_i^{\vec{j}}(z_i)|^2, \quad \text{with } S_i^{\vec{j}}(z_i) \equiv \langle \vec{j}, i | \vec{j}, n(z) \rangle. \quad (95)$$

From this expansion one concludes that $\|N(z_i)\|_{\vec{j}} \neq 0$ iff there exists at least one holomorphic section which does not vanish at z_i . Thus, the previous characterisation is equivalent to saying that $P_{\vec{j}}^s \equiv \text{SL}(2, \mathbb{C}) \cdot P_{\vec{j}}^{(0)}$ is the space of points, where at least one invariant holomorphic section does not vanish.

We can now use the analysis of the 4-valent sections performed in the previous section. From (76) one can infer that if $\vec{j}$ is regular (i.e. $j_1 \pm j_2 \pm j_3 \pm j_4 \neq 0$ for any choice of sign) then for any nonzero section $S^{\vec{j}}$, we have that $S^{\vec{j}}(z_i) = 0$ when $Z(z_i) = 0, 1, \infty$. On the other hand, it is clear that when $Z(z_i) \neq 0, 1, \infty$ (i.e. $z_i \neq z_j$) we can always find a section such that $S^{\vec{j}}(z_i) \neq 0$.

More precisely, it is immediate from (76) that $S^{\vec{j}}(z_i) \neq 0$ if $z_i \neq z_j$ for all nonzero sections if $\vec{j}$ is regular. On the other hand, if for instance $j_1 + j_2 > j_3 + j_4$, then $S^{\vec{j}}(z_i) \rightarrow 0$ when $z_{12} \rightarrow 0$ (that is, when $Z \rightarrow \infty$). Thus, what we have obtained is the characterisation of the orbit space as

$$P_{\vec{j}}^s \equiv \text{SL}(2, \mathbb{C}) \cdot P_{\vec{j}}^{(0)} = \mathbb{C} \setminus \{0, 1, \infty\} \times \text{SL}(2, \mathbb{C})$$

for regular¹² j_i .

¹¹ To be precise, Guillemin and Sternberg use a characterization in terms of non-vanishing invariant holomorphic sections, but this is equivalent to the non-vanishing of coherent intertwiners, as we see below.

¹² If j_i is not regular, then it is possible to find a section which is nonzero even for $Z = 0$ or 1 or ∞ , so that the set P^s is, in fact, bigger and includes also degenerate tetrahedra of zero volume. For instance, in the case where $j_1 + j_2 = j_3 + j_4$ there exists a section which is nonzero even when evaluated at $z_{12} = 0$ or $z_{34} = 0$.

C. The main correspondence

We now want to establish explicitly the isomorphism between the Hilbert space $H_{\vec{j}}$ which is the usual Hilbert space of $\text{SU}(2)$ intertwiners and the Hilbert space $H_{\vec{j}}^{(0)}$ obtained by first reducing and then quantizing. The latter is the span of coherent intertwiners satisfying the closure constraint. We impose the restriction that the set j_i is regular, i.e. $j_1 \pm j_2 \pm j_3 \pm j_4 \neq 0$ for any choice of sign. We start from the decomposition of the identity of $H_{\vec{j}}$, given by

$$\mathbb{1}_{\vec{j}} = \prod_i d_{j_i} \int_{P_{\vec{j}}} \prod_i d^2 N(z_i) ||\vec{j}, N(z_i)\rangle \langle \vec{j}, N(z_i)||. \quad (96)$$

The integral over $P_{\vec{j}}$ can be replaced by the integral of the stable set $P_{\vec{j}}^s$, since the complement is of measure zero. This integral can be further expanded as an integral over the constraint space and the $\text{SL}(2, \mathbb{C})$ orbit space due to the isomorphism

$$P_{\vec{j}}^s = P_{\vec{j}}^{(0)} / \text{SU}(2) \times \text{SL}(2, \mathbb{C}) = P_{\vec{j}}^{(0)} \times \text{SL}(2, \mathbb{C}) / \text{SU}(2)$$

proven in the previous section. In the following, we will view the space $\text{SL}(2, \mathbb{C}) / \text{SU}(2) \equiv H_3$ as the space¹³ of positive hermitian two by two matrices of determinant 1—that is, as a subspace of $\text{SL}(2, \mathbb{C})$. This space has a unique invariant measure and we normalize this measure to be

$$de^{iX} = \left(\frac{2}{|X|} \sinh \frac{|X|}{2} \right)^2 \prod_a dX^a \quad \text{with} \quad X = X^a \tau_a \in \mathfrak{su}(2), \quad |X|^2 = X^a X_a.$$

Taking advantage of this decomposition we can now show that

$$\int_{P_{\vec{j}}^s} \prod_i d^2 N_i f(N_i) = \int_{P_{\vec{j}}^{(0)}} d\mu^{(0)}(N_i) \det(G(N_i)) \left[\int_{H_3} \prod_i |\rho_h(N_i)|^2 f(h \cdot N_i) dh \right], \quad (97)$$

where $d\mu^{(0)}(N_i)$ is the $\text{SU}(2)$ invariant measure on $P_{\vec{j}}^{(0)}$:

$$d\mu^{(0)}(N_i) \equiv \prod_i d^2 N_i \delta^{(3)} \left(\sum_i j_i N_i \right),$$

and $\det(G(N_i))$ is the determinant of the metric (86) which has been computed in [17]:

$$\det(G(N_i)) = \left(\sum_i j_i \right) \sum_{i>j} j_i j_j |N_i, N_j|^2 - \frac{1}{6} \sum_{i,j,k} j_i j_j j_k ([N_i, N_j] | N_k)^2 \quad (98)$$

$$= \frac{(\sum_i j_i) V^2(L)}{j_1 j_2 j_3 j_4} \left(\sum_{i>j} j_i j_j |L^{ij}|^2 - V^2(L) \right). \quad (99)$$

¹³ The isomorphism between $\text{SL}(2, \mathbb{C}) / \text{SU}(2)$ and H_3 is given by $g \rightarrow h^2 = gg^\dagger$. This space is also the 3-dimensional Hyperbolic space.

The second equality is only valid if the N_i 's satisfy the closure constraint. In this equality we expressed the determinant in terms of the edge lengths L^{ij} (see eq. (32)) and the volume $V(L)$ of the tetrahedron¹⁴ using the relations established in section II B. It is interesting to note that the degenerate configurations are suppressed by the presence of this determinant.

In order to derive this integration formula we first recall that $G(X, Y)(N_i) = J(\widehat{X})H_Y(N_i)$. Thus, we have the identity

$$1 = \det(G(N_i)) \int_{H_3} dh \delta^{(3)}(h \cdot H) = \det(G(N_i)) \int_{H_3} dh \delta^{(3)}\left(h \cdot \left(\sum_i j_i N_i\right)\right). \quad (100)$$

Inserting this identity in the LHS of (97) and performing a change of variables $N_i \rightarrow h \cdot N_i$, one gets the RHS, since

$$d^2(h \cdot N(z)) = |\rho_h(z)|^2 d^2 N(z). \quad (101)$$

This proof is similar to the Faddeev–Popov proof of gauge fixing.

Given this lemma and the invariance property (81) of the state one arrives at

$$\mathbb{1}_{\vec{j}} = \prod_i d_{j_i} \int_{P_{\vec{j}}^{(0)}} d\mu_{\vec{j}}(N_i) \|\vec{j}, N_i\rangle \langle \vec{j}, N_i|, \quad (102)$$

where the “quantum” integration measure $d\mu(N_i)$ differs from the “classical” integration measure $\mu^{(0)}(N_i)$ by a factor which is $\text{SL}(2, \mathbb{C})$ invariant and dependent on $\vec{j}$:

$$d\mu_{\vec{j}}(N_i) \equiv d\mu^{(0)}(N_i) \det(G(N_i)) \left[\int_{H_3} \prod_i |\rho_h(z_i)|^{2(j_i+1)} dh \right] \quad (103)$$

What is remarkable about this identity is the fact that the integration is only over the constraint surface or the set of tetrahedra. This shows that coherent states which satisfy the closure constraint provide a basis of $H_{\vec{j}} = (V^{j_1} \dots \otimes V^{j_4})^{\text{SU}(2)}$. Since their labels correspond to classical tetrahedra, this means that these states have a very clear geometrical interpretation as tetrahedra with fixed areas and no longer need any interpretation as fuzzy tetrahedra, as was the case in the works of Barbieri [13] and Livine–Speziale [17].

We finally want to establish the asymptotic property of the measure term when the spins are uniformly rescaled, i.e. $j_i \rightarrow \lambda j_i$. The measure (103) involves an integral of a weight which is always smaller than one. In the large λ limit the integral therefore localises on its maximum point [32]. Since the N_i 's meet the constraint, the maximum is at $h = 1$. At this point the value of the integrand is equal to one and the integral is just determined asymptotically by the determinant of the matrix of quadratic fluctuation around the maximum. As we have established in eq. (85), this matrix is given by G , that is, $\sum_i 2j_i \ln |\rho_{e^{ix}}(z_i)| = -\lambda G(X, X) + O(X^3)$, therefore

$$d\mu_{\vec{j}}(N_i) \sim d\mu^{(0)}(N_i) \left(\frac{\pi}{\lambda}\right)^{\frac{3}{2}} \frac{1}{\sqrt{\det(G(N_i))}}. \quad (104)$$

¹⁴ More precisely, $V(L) \equiv \det(L^{14}, L^{24}, L^{34})$ is 6 times the volume of the tetrahedron.

VI. FK MODEL AND ASYMPTOTICS FOR A SINGLE 4-SIMPLEX

A. FK γ model

In this section, we will apply our central result, eq. (97), to the FK γ spin foam model [2]. For $\gamma < 1$, this model is identical to the EPRL model [4].

The model is defined on a two dimensional “spine” $\mathcal{S}$ which is the 2-dimensional skeleton of the intersection of Δ^* with Δ , where Δ is a simplicial complex (for more details, see ref. [2] and [7]). We denote by v , e and f the edges vertices, edges and faces of the dual complex Δ^* . We assign $SU(2) \times SU(2)$ group elements $\mathbf{g}_{ev} = (g_{ev}^+, g_{ev}^-)$ to each oriented pair ev , with the convention that $g_{ve} = g_{ev}^{-1}$, a point of S^2 denoted N_{ef} to every pair ef , and a pair of spins $j^\pm = |\gamma^\pm|j_f$ to every face f ¹⁵. A pair (vf) is called a wedge and it is the meeting point of two edges e, e' . The partition function of the model is given by

$$Z = \sum_{j_f} \int \prod_{f, e \subset f} d_{j_f^+} d_{j_f^-} dN_{ef} \prod_{v, e \supset v} d\mathbf{g}_{ev} \prod_{v, f \supset v} A_{vf}(j_f^\pm, \mathbf{g}_{ev}, N_{ef}), \quad (106)$$

where for $\gamma < 1$ the amplitude of the wedge vf is¹⁶

$$A_{vf}(j_f^\pm, \mathbf{g}_{ev}, N_{ef}) = \langle j_f^+, N_{ef} | T^{j_f^+}(g_{ev}^+ g_{ve'}^+) | j_f^+, N_{ef} \rangle \langle j_f^-, N_{ef} | T^{j_f^-}(g_{ev}^- g_{ve'}^-) | j_f^-, N_{ef} \rangle. \quad (108)$$

This amplitude can be described as a contraction of tensor products of coherent intertwiners from each edge: that is, if one denotes $\vec{j}_e \equiv (j_f)_{f \subset e}$ and $\vec{N}_e \equiv (N_{ef})_{f \subset e}$, one can obtain Z by contracting the invariant tensor

$$\bigotimes_e \int \prod_{f \supset e} dN_{ef} \left(\|\vec{j}_e^+, \vec{N}_e\| \langle \vec{j}_e^+, \vec{N}_e \| \otimes \|\vec{j}_e^-, \vec{N}_e\| \langle \vec{j}_e^-, \vec{N}_e \| \right). \quad (109)$$

When viewed in this way, it becomes clear that we can apply the main result proven in the previous section: first one replaces the integral over unconstrained N_{ef} by an integral over $P_{\vec{j}}^{(0)} \times H_3$, and then one uses the $SL(2, \mathbb{C})$ covariance in order to integrate out the dependence on the “fiber” H_3 . Thus, we get the amplitude

$$Z = \sum_{j_f} \prod_e \int_{P_{\vec{j}}^{(0)}} d\mu_{\vec{j}_e}^{\vec{j}_e}(N_{ef}) \int \prod_{v, e \supset v} d\mathbf{g}_{ev} \prod_{v, f \supset v} A_{vf}(j_f^\pm, \mathbf{g}_{ev}, N_{ef}), \quad (110)$$

¹⁵ $\gamma^\pm$ are determined by the Immirzi parameter γ as the smallest integers such that [5]

$$\frac{\gamma_+}{\gamma_-} = \frac{\gamma + 1}{\gamma - 1}, \quad \gamma^+ > 0. \quad (105)$$

¹⁶ The corresponding amplitude for the case $\gamma > 1$ would be

$$A_{vf}(j_f^\pm, \mathbf{g}_{ev}, N_{ef}) = \langle j_f^+, N_{ef} | T^{j_f^+}(g_{ev}^+ g_{ve'}^+) | j_f^+, N_{ef} \rangle \overline{\langle j_f^-, N_{ef} | T^{j_f^-}(g_{ev}^- g_{ve'}^-) | j_f^-, N_{ef} \rangle}, \quad (107)$$

with the overline denoting complex conjugation. For sake of brevity, we only treat the case $\gamma < 1$ here. The derivation and results for $\gamma > 1$ are essentially the same up to minor technical modifications.

where the new measure for N_{ef} , $f \supset e$, is

$$d\mu_e^{\vec{j}_e}(N_{ef}) \equiv \left(\prod_{f \supset e} d_{j_f^+} d_{j_f^-} d^2 N_{ef} \right) \delta^{(3)} \left(\sum_{f \supset e} j_f N_{ef} \right) \det(G_{\vec{j}_e}(N_{ef})) \left[\int_{H_3} \prod_{f \supset e} |\rho_h(z_{ef})|^{2(j_f^+ + j_f^- + 1)} dh \right].$$

The key point is that now we only integrate over vectors that satisfy the closure constraint, i.e. $\sum_{f \supset e} j_f N_{ef} = 0$ for each edge e .

So far we have only described the amplitude for a closed 2d spine $\mathcal{S}$, but it is straightforward to generalize it to an amplitude with boundary states. Such an extension is needed in the computation of the graviton propagator (see for example [33, 34, 35]).

It is known that in the case $\gamma < 1$ the boundary states are $SU(2)$ spin networks [5, 6]. Such states are characterised by a choice of a 4-valent graph Γ , which is the boundary of the spine $\mathcal{S}$, a choice of $SU(2)$ representations $k_{\bar{e}}$ associated with the edges $\bar{e}$ of Γ and the choice of an intertwiner $i_{\bar{v}} \in (\otimes_{\bar{e} \supset \bar{v}} V_{k_{\bar{e}}})^{SU(2)}$ for each vertex $\bar{v}$ of Γ . $i_{\bar{v}}$ is a label representing either a usual spin network intertwiner (in which case it is a half-integer) or a coherent intertwiner (in which case it is given by four unit vectors). The edge label $k_{\bar{e}}$ of Γ is related to the face label j_f of a face intersecting the boundary: namely, $k_{\bar{e}} = j_f^+ + j_f^-$ or $j_f = k_{\bar{e}}/(\gamma^+ - \gamma^-)$ (since $\gamma^- < 0$ for $\gamma < 1$).

In order to define the amplitude we just have to distinguish between the bulk edges e of the spine and the boundary edges of the spine which are in one-to-one correspondence with boundary vertices $\bar{v}$. Similarly, we distinguish the bulk faces f from boundary faces which are in one-to-one correspondence with boundary edges $\bar{e}$. Given these boundary data, we can specify the boundary amplitude as

$$Z(\Gamma, k_{\bar{e}}, i_{\bar{v}}) = \int \prod_{\bar{v}, \bar{e} \supset \bar{v}} d_{k_{\bar{e}}} dN_{\bar{v}\bar{e}} Z(j_{\bar{e}}, N_{\bar{v}\bar{e}}) \prod_{\bar{v}} \langle \vec{k}_{\bar{v}}, \vec{N}_{\bar{v}} \| \vec{k}_{\bar{v}}, i_{\bar{v}} \rangle. \quad (111)$$

Here, $j_{\bar{e}} = k_{\bar{e}}/(\gamma^+ - \gamma^-)$, and $Z(j_{\bar{e}}, N_{\bar{v}\bar{e}})$ is defined like the partition function (110) except that the integration and summation extends only over bulk variables:

$$Z(j_{\bar{e}}, N_{\bar{v}\bar{e}}) = \sum_{j_f} \int \prod_{f, e \subset f} d_{j_f^+} d_{j_f^-} dN_{ef} \prod_{v, e \supset v} d\mathbf{g}_{ev} \prod_{v, f \supset v} A_{vf}(j_f^\pm, \mathbf{g}_{ev}, N_{ef}). \quad (112)$$

One can show that this definition of the boundary amplitude agrees with the one given in [7]¹⁷. As before, we can apply the result of the previous section and replace the integral in (111) by an integral with measure μ where the closure constraint is imposed strongly:

$$Z(\Gamma, k_{\bar{e}}, i_{\bar{v}}) = \prod_{\bar{v}} \int_{P_j^{(0)}} d\mu_{\bar{v}}^{\vec{j}_v}(\vec{N}_{\bar{v}}) Z(j_{\bar{e}}, N_{\bar{v}\bar{e}}) \prod_{\bar{v}} \langle \vec{k}_{\bar{v}}, \vec{N}_{\bar{v}} \| \vec{k}_{\bar{v}}, i_{\bar{v}} \rangle \quad (114)$$

¹⁷ For this one uses the coherence property of coherent states, that is, the identity

$$|j^+, N\rangle \otimes |j^-, N\rangle = |k, N\rangle \quad \text{for } k = j^+ + j^-. \quad (113)$$




Figure 1: Dual of a 4-simplex with boundary: the boundary of the 4-simplex intersects dual edges at their center. The intersection points provide the base point for boundary data $n_{e_i f_{ij}}, u_{e_i}$.

with

$$d\mu_{\vec{v}}^{\vec{j}_{\vec{v}}}(\vec{N}_{\vec{v}}) \equiv \left(\prod_{\vec{v}, \vec{e} \supset \vec{v}} d_{k_{\vec{e}}} d^2 N_{\vec{v}\vec{e}} \right) \delta^{(3)} \left(\sum_{\vec{e} \supset \vec{v}} k_{\vec{e}} N_{\vec{v}\vec{e}} \right) \det(G_{\vec{j}_{\vec{v}}}(N_{\vec{v}\vec{e}})) \left[\int_{H_3} \prod_{\vec{e} \supset \vec{v}} |\rho_h(z_{\vec{v}\vec{e}})|^{2(k_{\vec{e}}+1)} dh \right] \quad (115)$$

and $k_{\vec{e}} = j_{\vec{e}}^+ + j_{\vec{e}}^-$.

B. Asymptotics of vertex amplitude

We now want to illustrate that the techniques used in [5] can be naturally extended to analyse the asymptotics of boundary amplitudes. Here, we will focus on the case of one 4-simplex (i.e. the vertex amplitude) reserving the general study for the future.

To describe the contribution of a single 4-simplex dual to a vertex v we number the edges at the vertex e_i , $i = 1, \dots, 5$ (see Fig. 1). Moreover, it will be convenient to abbreviate the notation and set $\mathbf{g}_i \equiv \mathbf{g}_{ve_i}$, $n_{ij} \equiv n_{e_i f_{ij}}$, $u_i \equiv u_{e_i}$, $j_{ij}^{\pm} \equiv j_{f_{ij}}^{\pm}$ etc. Then,

$$A_v(j_{ij}^{\pm}, N_{ij}) = \int \prod_i dg_i^+ dg_i^- \prod_{i < j} A_{ij}(j_{ij}^{\pm}, \mathbf{g}_i, N_{ij}) \quad (116)$$

where

$$A_{ij}(j_{ij}^{\pm}, \mathbf{g}_i, N_{ij}) = \langle j_{ij}^+, N_{ij} | T_{ij}^{j_{ij}^+} \left((g_i^+)^{-1} g_j^+ \right) | j_{ij}^+, N_{ji} \rangle \langle j_{ij}^-, N_{ij} | T_{ij}^{j_{ij}^-} \left((g_i^-)^{-1} g_j^- \right) | j_{ij}^-, N_{ji} \rangle. \quad (117)$$

We assume that the closure constraint is imposed, since, as shown in the previous section, we only have to integrate over constrained configurations in order to define the boundary state amplitude. Therefore,

$$\sum_{j: j \neq i} j_{ij} N_{ij} = 0. \quad (118)$$

Also, in order to use the results of [5] we need to assume that the integral is only over nondegenerate configurations, i.e. we impose in the integral the restriction

$$\det(g_1 \hat{U}, \dots, g_4 \hat{U}) \neq 0, \quad (119)$$

where $\hat{U} = (1, 0, 0, 0)^T$ is a reference vector. We call the restricted nondegenerate amplitude A_v^{nd} .

Before we can state our result on the asymptotics, we need to introduce a number of definitions. Recall from refs. [5, 7] that the variables $j_f^\pm$ and N_{ef} define a bivector X_{ef}^γ that represents the discrete analogue of the B -field. Its selfdual and anti-selfdual components are

$$\mathbf{X}_{ef}^\gamma = (j_f^+ N_{ef}, j_f^- N_{ef}), \quad N_{ef} \equiv N_{ef}^i \sigma_i. \quad (120)$$

It is convenient to introduce a *simple* bivector X_{ef} by setting

$$X^{\gamma^\pm} \equiv \gamma^\pm X^\pm, \quad \text{so that} \quad \mathbf{X}_{ef} = (j_f N_{ef}, j_f N_{ef}), \quad (121)$$

and

$$X_{ef}^\gamma = \frac{1}{2} (\gamma^+ + \gamma^-) \left(X_{ef} + \frac{1}{\gamma} \star X_{ef} \right). \quad (122)$$

From lemma C.1. in [7] we also know that this corresponds to a bivector

$$X_{ef} = j_f \hat{U} \wedge \hat{N}_{ef}, \quad (123)$$

where $\hat{U} = (1, 0, 0, 0)^T$ and $\hat{N}_{ef}$ is the 4-vector $\hat{N}_{ef} = (0, N_{ef})$ and N_{ef} the previously defined 3-vector. In terms of our index notation, this means that

$$X_{ij} = j_{ij} \hat{U} \wedge \hat{N}_{ij}. \quad (124)$$

Let us next define vectors

$$V_{ij} = R(n_{ij}, n_{ij})(0, 1, 0, 0)^T, \quad W_{ij} = R(n_{ij}, n_{ij})(0, 0, 1, 0)^T, \quad (125)$$

where $R(g^+, g^-)$ denotes the projection of the $\text{SU}(2) \times \text{SU}(2)$ element $\mathbf{g} = (g^+, g^-)$ to $\text{SO}(4)$. We will see in appendix A that V_{ij} and W_{ij} are vectors in the plane orthogonal to X_{ij} . Given the vectors $\hat{U}$, $\hat{N}_{ij}$, V_{ij} and W_{ij} , we specify angles θ_{ij} and $\tilde{\theta}_{ij}$ by

$$\cos \theta_{ij} = \hat{U} \cdot g_{ij} \hat{U}, \quad \sin \theta_{ij} = \hat{U} \cdot g_{ij} \hat{N}_{ji}, \quad (126)$$

$$\cos \tilde{\theta}_{ij} = V_{ij} \cdot g_{ij} V_{ji}, \quad \sin \tilde{\theta}_{ij} = V_{ij} \cdot g_{ij} W_{ji}, \quad (127)$$

where $g\hat{U}$, $g\hat{N}$ etc. stands for the action of the $\text{SO}(4)$ element g on 4-vectors. These angles are needed to express the asymptotics of the vertex amplitude.

The asymptotics is the result of a saddle point approximation of the integral over $\mathbf{g}_i$ in (116). Since

$$|\langle j^+ N_i | g_i^{-1} g_j | j^+ N_j \rangle| = \left| \frac{1}{2} (1 + (g_i N_i g_i^{-1} | g_j N_j g_j^{-1})) \right|^{j^+} \leq 1, \quad (128)$$

with the equality satisfied iff $g_i N_i g_i^{-1} = g_j N_j g_j^{-1}$, it follows that the saddle points (the points where the integrand is maximal) are determined by the equations

$$g_i X_{ij} g_i^{-1} = g_j X_{ji} g_j^{-1}. \quad (129)$$

This equation should be supplemented by the closure constraint (118) of the boundary data, which can be written as

$$\sum_{j \neq i} X_{ij} = 0. \quad (130)$$

When a solution exists and nondegeneracy is assumed, these equations imply that there is a 4-simplex at the vertex v which is consistent with the tetrahedra in the boundary¹⁸. This follows from a straightforward extension of proposition VII.1, [5] to a complex with boundary. The difference to the analysis in [5] is only that the boundary conditions for the 4-simplex are now fixed by the boundary tetrahedra, and not by neighbouring 4-simplices.

The geometry of this 4-simplex is determined by the 4-simplex tetrad U_i (see def. VI.3 and prop. VI.4 of [5]), i.e. a set of five 4d vectors which satisfy the closure condition $\sum_i U_i = 0$ and whose direction is, up to sign, given by $g_i \hat{U}$:

$$\frac{U_i}{|U_i|} = \pm g_i \hat{U}. \quad (131)$$

The bivectors X_{ij} arise from wedge products of the tetrad vectors:

$$g_i X_{ij} g_i^{-1} = \epsilon V_4(v)(U_i \wedge U_j). \quad (132)$$

ϵ is a sign factor and determined by the boundary data. The group elements g_i are, up to sign, equal to the spin connection, which maps the vertex geometry of the tetrahedra determined by U_i to the boundary geometry of the tetrahedra determined by N_{ij} . More precisely, given the U_i , we can compute the geometry of any of the five tetrahedra of the 4-simplex. As a result of the stationary phase equation (129), this geometry is the same as the one derived directly from the boundary data (see sec. II B). They only differ by a rotation of frame specified by the spin connection. These results follow from proposition VII.1 and proposition B.2 of [5].

From equality (131) and definition (126) one sees that the angles θ_{ij} are the dihedral angles of the 4-simplex geometry determined by U_i modulo π . The subtlety is that $g_i \hat{U}$ can differ from the normalised tetrad either by a sign or by a global reflection¹⁹ around $\hat{U}$, and thus θ_{ij} equals the dihedral angle only up to π or up to a global sign. The reflection around $\hat{U}$ does not affect the $\tilde{\theta}$ angles, since they correspond to rotations in a plane orthogonal to $\hat{U}$.

Here, we will restrict our analysis to the case where the spins $j_{ij}^\pm$ are integers. In this case, the difference between θ_{ij} and $\theta_{ij} + \pi$ is of no consequence in the asymptotics (see eq. (133) below).

Based on the saddle point approximation we can now deduce the asymptotics of the integral (116). When the eqns. (129) have a solution, we find that

$$A_v^{\text{nd}} \sim \frac{1}{(4\pi)^4} \frac{1}{\sqrt{\det(G_+) \det(G_-)}} \exp \left[-i \sum_{i < j} \left((\gamma^+ + \gamma^-) j_{ij} \theta_{ij} + (\gamma^+ - \gamma^-) j_{ij} \tilde{\theta}_{ij} \right) \right] + (\theta \rightarrow -\theta), \quad (133)$$

¹⁸ See [36] for a canonical analysis of the same equations.

¹⁹ This last point was initially overlooked and corrected thanks to reference [25].

where the second term is obtained by the reflection $(\theta, \tilde{\theta}) \rightarrow (-\theta, \tilde{\theta})$. When there are no nondegenerate solutions, the amplitude is exponentially suppressed. In the latter case, there is no nondegenerate 4-simplex which is consistent with the boundary tetrahedra.

Since $\mathcal{A}_{ij} = (\gamma^+ + \gamma^-)j_{ij}/2$ is the area of the triangle dual to the face ij and θ_{ij} is the dihedral angle, the first term in the exponential is the 4d Regge action of the 4-simplex. The second term depends on the additional angles $\tilde{\theta}_{ij}$. One can show (directly or using the results of [5]) that these terms drop out when the saddle point approximation is applied to a bulk region consisting of several 4-simplices. More precisely, for a face f in the bulk $\sum_{f \supset e} \tilde{\theta}_{ef} = 0$ modulo π . Therefore, eq. (133) is consistent with the bulk asymptotics derived in our previous paper [5]. It can be also understood that the extra angle $\tilde{\theta}_{ij}$ is essentially due to a $U(1)$ gauge symmetry of coherent states. Indeed, we have labelled coherent states so far by elements $N \in \text{SU}(2)/U(1)$ and corresponding group elements $n \in \text{SU}(2)$. We could label them just as well by $\text{SU}(2)$ elements $n' = n e^{i\frac{\phi}{2}\sigma_3}$. If we do so, we see that the vertex amplitude becomes²⁰

$$A_v(n'_{ij}) = A_v(n_{ij}) e^{i(\gamma^+ - \gamma^-)j_{ij} \sum_{i < j} (\phi_{ij} - \phi_{ji})} . \quad (134)$$

Thus, the angles $\tilde{\theta}_{ij}$ can be locally gauged to zero.

The determinants $\det(G_{\pm})$ are determinants of 4×3 by 4×3 matrices: the rows and columns are labelled by pairs ai , where i denotes four tetrahedra of the 4-simplex, and a is a 3-dimensional index labelling a basis of $\mathfrak{su}(2)$. The matrix elements of $G_{\pm}$ are sums of the hermitian matrices (56):

$$G_{\pm}^{ia,ib} = \sum_{j:j \neq i} j_{ij}^{\pm} (\delta^{ab} - N_{\pm ij}^a N_{\pm ij}^b) , \quad (135)$$

$$G_{\pm}^{ia,jb} = -j_{ij}^{\pm} (\delta^{ab} - N_{\pm ij}^a N_{\pm ij}^b - i\epsilon^{abc} N_{\pm ij}^c(z)) , \quad i \neq j , \quad (136)$$

with $N_{\pm ij} \equiv g_i^{\pm} N_{ij} (g_i^{\pm})^{-1}$. A detailed derivation of formula (133) is provided in appendix A.

Eq. (133) describes the asymptotics of the vertex amplitude A_v^{nd} , in which the integration over g_i is constrained by the nondegeneracy condition (119). Therefore, there can only appear saddle points corresponding to nondegenerate 4-simplices. For these points, our formula agrees, up to numerical factors, with the asymptotics obtained in ref. [25]. In the latter work the vertex amplitude is treated without any restriction, so there appear additional terms corresponding to degenerate configurations.

VII. SUMMARY AND DISCUSSION

Let us summarize the contents and results of this paper: we followed a reduction-before-quantization approach to construct the Hilbert space of a tetrahedron in $\mathbb{R}^3$. Guided by theorems of Guillemin & Sternberg and Hall [19, 38] we related this quantization to the more conventional Dirac quantization, where one quantizes first and imposes the constraints afterwards. We started by describing the classical phase space of tetrahedra and the Guillemin-Sternberg theorem at the classical level (relating the phase spaces $P^{(0)}/\text{SU}(2)$

²⁰ Observe that for $\gamma < 1$ we have that $\gamma^- < 0$.

and $P^s/\mathrm{SL}(2, \mathbb{C})$). Then, we reviewed properties of group coherent states which we need to connect classical phase space and quantum theory. Section IV outlined the content of the Guillemin–Sternberg isomorphism at the quantum level. We then gave a proof and explicit construction of this map for the quantum tetrahedron (sec. V): the central formula is eq. (102), which expresses the resolution of identity by an integral over coherent states labelled by tetrahedra. The underlying idea is the following: coherent states provide an overcomplete set of states and they are labelled by phase space variables. By Guillemin & Sternberg’s theorem the unconstrained phase space is naturally fibered over the constraint phase space with the fiber being isomorphic to the imaginary part of the gauge group. Therefore, we can express the integral over coherent states as an integral over the constrained variables and an integral over the hermitian elements of $\mathrm{SL}(2, \mathbb{C})$. Due to the $\mathrm{SL}(2, \mathbb{C})$ invariance of the state the fiber integral decouples via a Faddeev–Popov trick and gives an explicit measure factor.

In the last section, this result was applied to the $\mathrm{FK}\gamma$ model in order to express the path integral in terms of tetrahedral variables²¹. Using this, we determined the asymptotics of the vertex amplitude in the nondegenerate sector. Up to numerical factors, our result coincides with the nondegenerate contribution in the asymptotic formula by Barrett et al. [25].

A main motivation for this work came from a previous result, where we showed that in the semiclassical limit spin foam amplitudes reduce to the Regge action of discrete geometries [5]. On the one hand, this showed a clear link between the spin foam model and discretized general relativity. On the other hand, it also made very clear that, a priori, spin foams are not geometries. The spin foam model can be seen as a first–order path integral, where the “metric” variables are not tetrads, but more general objects, that only become tetrads when we go on–shell w.r.t. the reality of the action and the equations of motion for the connection (which includes the closure constraint).

For this reason, it is important to know how the amplitudes behave as one moves away from the constraint surface. When computing a graviton propagator, for example, one has to integrate over *all* fluctuations around a background, which includes those that do not fulfill the constraints. Thus, we were interested in expressing the spin foam amplitude as a function of an on–shell background plus something else which would parametrize the points away from the constraint surface.

With equation (102) we have obtained precisely such a formula for the closure constraint. The configurations are labelled by geometrical tetrahedra and $\mathrm{SL}(2, \mathbb{C})$ elements: the former characterize the point on the constraint surface from which we start to generate an arbitrary point by application of $\mathrm{SL}(2, \mathbb{C})$. Our formula tells us how the amplitude changes as a function of the $\mathrm{SL}(2, \mathbb{C})$ element, or equivalently, as a function of the distance from the constraint surface.

This also answers another question that motivated our work: in [5] an essential, but somewhat mysterious ingredient was the fact that the action possessed an imaginary part²². This imaginary part was absolutely essential in order to show that the amplitude had the right semiclassical limit: the condition that the imaginary part vanishes supplemented the equations of motion and ensured that only geometrical configurations dominate. We now see that this imaginary part is related to the existence of a natural metric on the Guillemin–Sternberg

²¹ For $\gamma < 1$, this applies also to the EPRL model [4], since in this case the FK and EPRL model are the same.

²² By the action we mean here S such that $A = e^{iS}$, where A is the amplitude.

fibration of phase space. The latter measures how far we are from “metricity”. Remarkably, as we have seen, some of the non-metrical dependence can be explicitly integrated out.

Our result is a further step in the direction initiated by Livine and Speziale’s coherent state method: namely, the attempt to express quantum states in terms of geometrical quantities. This approach has already led to considerable progress and clarification in the construction of spin foam models. With the new overcomplete set of states defined here, we go yet another step in this direction: now the states are not only labelled by spins and normal vectors, but, in addition, these normal vectors close, so they truly define a geometrical tetrahedron. In this sense, these states are closer to the notion of geometry than the spin network intertwiners, and one is more justified to call them states of quantum geometry.

We believe that the use of such states could have various advantages in spin foam models and also in canonical loop quantum gravity. It could give access to perturbation theory and Feynman diagrams, since the amplitude is now conveniently expressed as a function of the on-shell background and an “off-shell parameter”. Thus, the path integral resembles a lattice path integral over a tetrad field, a connection and further parameters, and it seems conceivable that one can derive Feynman diagrams in a similar way as in lattice gauge theory.

The methods developed in our paper should also shed new light on the coherent states of canonical loop quantum gravity [37]. A notable difference between our states and the ones described there is the fact that the latter states are designed to give semiclassical peakedness in both the connection and area, while our states are peaked on explicit tetrahedral geometries. It would be interesting to understand the relationship between the two approaches.

The LQG coherent states arise in the definition of the coherent state transform $L^2(\mathcal{A}) \rightarrow L^2_{\text{hol}}(\mathcal{A}^{\mathbb{C}})$ [37] and extend Hall’s notion of group coherent states [38]. In ref. [39], Bahr and Thiemann have applied group averaging on these states, and obtained gauge-invariant coherent states which are the analog of the Livine–Speziale coherent states. Such states are not only invariant under the gauge group $\mathcal{G}$, but also under the complexified group $\mathcal{G}^{\mathbb{C}}$, as in our case. What is missing so far in the analysis of these states (and this would constitute the analog of what we show here) is the proof that the Hilbert space of gauge invariant states is in fact unitarily equivalent to the Hilbert space of holomorphic functionals $L^2_{\text{hol}}(\mathcal{A}^s/\mathcal{G}^{\mathbb{C}})$ with an appropriate measure, where $\mathcal{A}^s$ is the analog of the semistable set P^s_j . Such an isomorphism would imply that all gauge invariant coherent states are properly peaked and it would also provide an explicit parametrisation and scalar product on the space of coherent states on which the Gauss law is imposed. We hope to come back to these issues in the future by expanding on the techniques that we have developed here.

In summary: our work continues in the spirit of recent developments that emphasize the geometric aspect of spin foam models and loop quantum gravity. We are hopeful that this will lead to further progress in linking the quantum with the classical theory and the spin foam with the canonical approach, and that it will bring us closer to answering the decisive question: namely, if this theory of quantum gravity is, in fact, consistent with classical general relativity.

Acknowledgments

We thank E. Alesci, E. Livine, B. Dittrich and S. Speziale for discussions. Research at Perimeter Institute is supported by the Government of Canada through Industry Canada and by the Province of Ontario through the Ministry of Research & Innovation.

Appendix A: DERIVATION OF ASYMPTOTICS

In this appendix we give more details on the derivation of the asymptotics (133). We will first focus on the edges e_1 and e_2 and write $N_1 \equiv N_{12}$, $N_2 \equiv N_{21}$ and $j \equiv j_{12}$ for simplicity. The associated group elements are $n_1 \equiv n_{12}$ and $n_2 \equiv n_{21}$.

From the edges e_1 , e_2 and the face f_{12} we get the amplitude factor

$$A_{12} = \langle j^+ n_1 | (g_1^+)^{-1} g_2^+ | j^+ n_2 \rangle \langle j^- n_1 | (g_1^-)^{-1} g_2^- | j^- n_2 \rangle. \quad (\text{A1})$$

Let us set $n_2^\pm = (g_1^\pm)^{-1} g_2^\pm n_2$. Then,

$$\left| \langle j^+ n_1 | (g_1^+)^{-1} g_2^+ | j^+ n_2 \rangle \right| = \left| \langle j^+ N_1 | j^+ N_2^+ \rangle \right| = \left(\frac{1}{2} (1 + N_1 \cdot N_2^+) \right)^{j^+}, \quad (\text{A2})$$

$$\left| \langle j^- n_1 | (g_1^-)^{-1} g_2^- | j^- n_2 \rangle \right| = \left| \langle j^- N_1 | j^- N_2^- \rangle \right| = \left(\frac{1}{2} (1 + N_1 \cdot N_2^-) \right)^{j^-}, \quad (\text{A3})$$

where N_2^+ and N_2^- are the unit 3-vectors associated to the group elements n_2^+ and n_2^- . On the other hand,

$$\langle j^+ n_1 | (g_1^+)^{-1} g_2^+ | j^+ n_2 \rangle = \left(\langle \tfrac{1}{2} - \tfrac{1}{2} | n_1^{-1} (g_1^+)^{-1} g_2^+ n_2 | \tfrac{1}{2} - \tfrac{1}{2} \rangle \right)^{2j} = e^{ij^+ \phi^+} \cos^{2j^+} (\varphi^+/2), \quad (\text{A4})$$

$$\langle j^- n_1 | (g_1^-)^{-1} g_2^- | j^- n_2 \rangle = \left(\langle \tfrac{1}{2} - \tfrac{1}{2} | n_1^{-1} (g_1^-)^{-1} g_2^- n_2 | \tfrac{1}{2} - \tfrac{1}{2} \rangle \right)^{2j} = e^{ij^- \phi^-} \cos^{2j^-} (\varphi^-/2) \quad (\text{A5})$$

for some angles $\phi^\pm$ and $\varphi^\pm$.

In the large spin limit, the moduli (A2) and (A3) are exponentially suppressed unless $N_1 = N_2^+ = N_2^-$. This condition is equivalent to the statement that

$$X_{12} = g_1^{-1} g_2 X_{21} g_2^{-1} g_1. \quad (\text{A6})$$

If this is fulfilled, the moduli are 1, the angles $\varphi^\pm$ are zero, and there only remain the phases $e^{ij^\pm \phi^\pm}$. In this case,

$$n_1^{-1} (g_1^\pm)^{-1} g_2^\pm n_2 = e^{-i\phi^\pm J_3} \quad (\text{A7})$$

and

$$A_{12} = \langle j n_1 | g_1^{-1} g_2 | j n_2 \rangle \langle j^- n_1 | (g_1^-)^{-1} g_2^- | j^- n_2 \rangle = e^{i(j^+ \phi^+ + j^- \phi^-)}. \quad (\text{A8})$$

By the same token we obtain the equations

$$X_{ij} = g_i^{-1} g_j X_{ji} g_j^{-1} g_i. \quad (\text{A9})$$

for the other pairs i, j .

As explained in sec. VI B, if there is a solution to eqns. (A9), it implies the existence of a 4-simplex at v whose area bivectors are the bivectors $\epsilon \star (g_i X_{ij} g_i^{-1})$. The vectors $g_i \hat{U}$ provide the normal vectors of tetrahedra at the basepoint v . Consequently, the angle θ_{12} defined by

$$\cos \theta_{12} = \hat{U} \cdot g_{12} \hat{U}, \quad (\text{A10})$$

$$\sin \theta_{12} = \hat{U} \cdot g_{12} \hat{N}_2, \quad (\text{A11})$$

is the dihedral angle²³ θ_{12} between the tetrahedra corresponding to the edge e_1 and e_2 .

To evaluate the phase in (A8), it is convenient to view the group elements in (A7) as the result of a gauge transformation which sends n_1 and n_2 to the identity:

$$g'_{12} = n_1^{-1} (g_1^\pm)^{-1} g_2^\pm n_2, \quad (\text{A12})$$

$$n'_1 = \mathbb{1}, \quad (\text{A13})$$

$$n'_2 = \mathbb{1}. \quad (\text{A14})$$

By equation (A7) we have that $(g'_{12})^\pm = e^{-i\phi^\pm J_3}$. The homomorphism $h : \mathfrak{su}(2) \oplus \mathfrak{su}(2) \rightarrow \mathfrak{so}(4)$ maps $\mathfrak{su}(2) \oplus \mathfrak{su}(2)$ elements $X^+ + X^- = X^{+i}\sigma_i + X^{-i}\sigma_i$ into $\mathfrak{so}(4)$ elements

$$2X = h(X^+, X^-), \quad (\text{A15})$$

where

$$X_i^\pm = \frac{1}{2} \epsilon_{0i}{}^{jk} X_{jk} \pm X_{0i}. \quad (\text{A16})$$

Inversion of the last equation gives

$$X_{ij} = \frac{1}{2} \epsilon_i{}^{jk} (X_k^+ + X_k^-), \quad (\text{A17})$$

$$X_{0i} = \frac{1}{2} (X_i^+ - X_i^-). \quad (\text{A18})$$

In the case of the group element g'_{12} , we have $X^\pm = -\phi^\pm J_3$, so

$$X_{ij} = -\frac{1}{4}(\phi^+ + \phi^-)\epsilon_{ij3}, \quad (\text{A19})$$

$$X_{0i} = -\frac{1}{4}(\phi^+ - \phi^-)\delta_{i3}, \quad (\text{A20})$$

and

$$g'_{12} = e^{2X} = e^{-\frac{1}{2}(\phi^+ - \phi^-)J_{03}} e^{-\frac{1}{2}(\phi^+ + \phi^-)J_{12}}. \quad (\text{A21})$$

Thus, if we define the angles α and $\tilde{\alpha}$ by

$$g'_{12} = e^{\alpha J_{03}} e^{\tilde{\alpha} J_{12}}, \quad (\text{A22})$$

we find that

$$\alpha = -(\phi^+ - \phi^-)/2, \quad \tilde{\alpha} = -(\phi^+ + \phi^-)/2. \quad (\text{A23})$$

Therefore,

$$j^+\phi^+ + j^-\phi^- = -(j^+ - j^-)\alpha - (j^+ + j^-)\tilde{\alpha}. \quad (\text{A24})$$

Using that $j^\pm = |\gamma^\pm|j$ and $\gamma^- < 0$ for $\gamma < 1$ (see [5]), we then arrive at

$$j^+\phi^+ + j^-\phi^- = -(\gamma^+ + \gamma^-)j\alpha - (\gamma^+ - \gamma^-)j\tilde{\alpha}. \quad (\text{A25})$$

After the rotation (A12)–(A14), the 4-vectors $\hat{N}_1$ and $\hat{N}_2$ become $\hat{N}'_1 = \hat{N}'_2 = (0, 0, 0, 1)^T$ and the transformed bivectors equal

$$X'_{12} = j\hat{U} \wedge \hat{N}'_1 = jJ_{03} = X'_{21}. \quad (\text{A26})$$

²³ As we discussed in the main part, it is actually the dihedral angle modulo π , due to the signs in eq. (131).

Therefore,

$$\cos \alpha = \hat{U} \cdot g'_{12} \hat{U}, \quad (\text{A27})$$

$$\sin \alpha = \hat{U} \cdot g'_{12} \hat{N}'_2, \quad (\text{A28})$$

$$\cos \tilde{\alpha} = (0, 1, 0, 0)^T \cdot g'_{12} (0, 1, 0, 0)^T, \quad (\text{A29})$$

$$\sin \tilde{\alpha} = (0, 1, 0, 0)^T \cdot g'_{12} (0, 0, 1, 0)^T. \quad (\text{A30})$$

We know that $\hat{U}$ is not affected by the transformation (A12)–(A14)²⁴, and $\hat{N}$ goes from $\hat{N}$ to $\hat{N}'$. Hence, if we reverse this rotation, we obtain

$$\cos \alpha = \hat{U} \cdot g_{12} \hat{U}, \quad (\text{A31})$$

$$\sin \alpha = \hat{U} \cdot g_{12} \hat{N}_2, \quad (\text{A32})$$

$$\cos \tilde{\alpha} = R(n_1, n_1)(0, 1, 0, 0)^T \cdot g_{12} R(n_2, n_2)(0, 1, 0, 0)^T, \quad (\text{A33})$$

$$\sin \tilde{\alpha} = R(n_1, n_1)(0, 1, 0, 0)^T \cdot g_{12} R(n_2, n_2)(0, 0, 1, 0)^T, \quad (\text{A34})$$

where we recognize the vectors V_{ij} and W_{ij} on the right-hand side (see eqns. (127)). Since $(0, 1, 0, 0)^T$ and $(0, 0, 1, 0)^T$ are orthogonal to the plane X'_{12} , the vectors V_{12} and W_{12} are orthogonal to X_{12} .

We conclude that $\alpha = \theta_{12}$ and $\tilde{\alpha} = \tilde{\theta}_{12}$, and hence

$$j^+ \phi^+ + j^- \phi^- = -(\gamma^+ + \gamma^-) j \theta_{12} - (\gamma^+ - \gamma^-) j \tilde{\theta}_{12}. \quad (\text{A35})$$

Thus, we obtain that

$$A_{12} = \exp \left[-i(\gamma^+ + \gamma^-) j \theta_{12} - i(\gamma^+ - \gamma^-) j \tilde{\theta}_{12} \right], \quad (\text{A36})$$

at the saddle point, or more generally

$$A_{ij} = \exp \left[-i(\gamma^+ + \gamma^-) j \theta_{ij} - i(\gamma^+ - \gamma^-) j \tilde{\theta}_{ij} \right]. \quad (\text{A37})$$

After taking into account determinant factors from the saddle point approximation, this yields eq. (133). The determinant factors can be easily obtained by expanding the logarithm of the amplitude around the stationary phase and using repeatedly (54) and (56). These formula show that the action in the exponential is at quadratic order given by $S = -\frac{1}{4} G_{+i,j}^{a,b} X_a^{+i} X_b^{+j} - \frac{1}{4} G_{-i,j}^{a,b} X_a^{-i} X_b^{-j} + O(X^3)$ with $g_i^\pm = \exp(X_a^{\pm i} \tau^a) \Omega_i^\pm$ and $\Omega_i^\pm$ denotes the spin connection which solves the stationary equations. Eventually, we have a factor $(4\pi)^{12}$ coming from the stationary phase evaluation and a factor $(4\pi)^{-16}$ coming from the normalisation of the measure on $\text{SU}(2)$. Because of the overall gauge invariance $g_i^\pm \rightarrow g^\pm g_i^\pm$ there are 8 integrations over $\text{SU}(2)$ to be performed instead of 10. Each $\text{SU}(2)$ integration provides a factor $(4\pi)^{-2}$, because the normalised $\text{SU}(2)$ measure is

$$\frac{1}{(4\pi)^2} \left(\frac{2}{|X|} \sin \frac{|X|}{2} \right)^2 \prod_a dX^a, \quad |X| < 2\pi. \quad (\text{A38})$$

²⁴ This can be also checked directly by using the equation $g^- x^I \sigma_I^E (g^+)^{-1} = (R(\mathbf{g})x)^I \sigma_I^E$, which relates $\text{SU}(2) \times \text{SU}(2)$ to $\text{SO}(4)$ transformations.

-
- [1] J. Engle, R. Pereira, and C. Rovelli, *The loop-quantum-gravity vertex-amplitude*, *Phys. Rev. Lett.*, 99:161301, 2007, 0705.2388.
 - [2] L. Freidel and K. Krasnov, *A new spin foam model for 4d gravity*, *Class. Quant. Grav.* **25**, 125018 (2008) arXiv:0708.1595.
 - [3] E. R. Livine and S. Speziale, *Consistently Solving the Simplicity Constraints for Spinfoam Quantum Gravity*, *Europhys. Lett.* **81**, 50004 (2008) arXiv:0708.1915.
 - [4] J. Engle, E. Livine, R. Pereira, and C. Rovelli, *LQG vertex with finite Immirzi parameter*, *Nucl. Phys.*, B799:136–149, 2008, arXiv:0711.0146.
 - [5] F. Conrady and L. Freidel, “On the semiclassical limit of 4d spin foam models,” arXiv:0809.2280 [gr-qc].
 - [6] J. Engle, R. Pereira, and C. Rovelli, *Flipped spinfoam vertex and loop gravity*, *Nucl. Phys.*, B798:251–290, 2008, 0708.1236.
 - [7] F. Conrady and L. Freidel, “Path integral representation of spin foam models of 4d gravity,” *Class. Quant. Grav.* **25**, 245010 (2008) [arXiv:0806.4640 [gr-qc]].
 - [8] L. Freidel, K. Krasnov and R. Puzio, “BF description of higher-dimensional gravity theories,” *Adv. Theor. Math. Phys.* **3**, 1289 (1999) [arXiv:hep-th/9901069].
 - [9] J.W. Barrett and L. Crane, *Relativistic spin networks and quantum gravity*, *J. Math. Phys.*, 39:3296–3302, 1998, gr-qc/9709028.
 - [10] J. C. Baez and J. W. Barrett, “The quantum tetrahedron in 3 and 4 dimensions,” *Adv. Theor. Math. Phys.* **3**, 815 (1999) [arXiv:gr-qc/9903060].
 - [11] E. Buffenoir, M. Henneaux, K. Noui and Ph. Roche, “Hamiltonian analysis of Plebanski theory,” *Class. Quant. Grav.* **21**, 5203 (2004) [arXiv:gr-qc/0404041].
 - [12] A. Baratin, C. Flori and T. Thiemann, “The Holst Spin Foam Model via Cubulations,” arXiv:0812.4055 [gr-qc].
 - [13] A. Barbieri, “Quantum tetrahedra and simplicial spin networks,” *Nucl. Phys. B* **518**, 714 (1998) [arXiv:gr-qc/9707010].
 - [14] M. P. Reisenberger and C. Rovelli, “*Sum over surfaces* form of loop quantum gravity,” *Phys. Rev. D* **56**, 3490 (1997) [arXiv:gr-qc/9612035].
 - [15] R. De Pietri and L. Freidel, *so(4) Plebanski Action and Relativistic Spin Foam Model*, *Class. Quant. Grav.*, 16:2187–2196, 1999, gr-qc/9804071.
 - [16] M. P. Reisenberger, *Classical Euclidean general relativity from *left-handed area = right-handed area**, gr-qc/9804061.
 - [17] E. R. Livine and S. Speziale, “A new spinfoam vertex for quantum gravity,” *Phys. Rev. D* **76**, 084028 (2007) [arXiv:0705.0674 [gr-qc]].
 - [18] C. Rovelli and S. Speziale, “A semiclassical tetrahedron,” *Class. Quant. Grav.* **23**, 5861 (2006) [arXiv:gr-qc/0606074].
 - [19] V. Guillemin and S. Sternberg. Geometric quantization and multiplicities of group representations. *Invent. Math.*, 67(3):515–538, 1982.
 - [20] V. Guillemin and S. Sternberg. “Symplectic techniques in Physics”, Cambridge University Press, 1993.
 - [21] M. Blau, “Symplectic geometry and geometric quantisation”
<http://www.blau.itp.unibe.ch/Lecturenotes.html>.

- [22] Justin Roberts. Classical $6j$ -symbols and the tetrahedron. *Geom. Topol.*, 3:21–66 (electronic), 1999.
- [23] L. Charles “On the quantization of polygon spaces,” arXiv:0806.1585v1 [math.SG].
- [38] B. Hall, W. Kirwin “Unitarity in ”quantization commutes with reduction”” *Comm. Math. Phys.* 275 (2007), no. 3, pages 401 – 442.
- [25] J. W. Barrett, R. J. Dowdall, W. J. Fairbairn, H. Gomes and F. Hellmann, “Asymptotic analysis of the EPRL four-simplex amplitude,” arXiv:0902.1170 [gr-qc].
- [26] D. Mumford, J. Fogarty, F. Kirwan, “Geometric invariant theory”, 3ed, Springer, 1994.
- [27] Kapovich M. and Millson “The symplectic geometry of Polygons in Euclidian space” *J. Journal of Differential Geometry*, Vol. 44 (1996) p. 479-513.
- [28] H. Waelbroeck and J. A. Zapata, “2 + 1 Covariant Lattice Theory and t’Hooft’s Formulation,” *Class. Quant. Grav.* **13**, 1761 (1996) [arXiv:gr-qc/9601011].
- [29] Z. Kadar, “Polygon model from first order gravity,” *Class. Quant. Grav.* **22**, 809 (2005) [arXiv:gr-qc/0410012].
- [30] A. Perelomov, “Generalized coherent states and their applications”, Springer-Verlag, 1986.
- [31] N. Vilenkin and A. Klimyk, “Representations of Lie groups and special functions” Kluwer academic Publisher, Vol. 1.
- [32] Lars Hörmander. *The analysis of linear partial differential operators. I*, volume 256 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, second edition, 1990. Distribution theory and Fourier analysis.
- [33] L. Modesto and C. Rovelli. *Particle scattering in loop quantum gravity*, *Phys. Rev. Lett.*, 95:191301, 2005, gr-qc/0502036.
- [34] E. Alesci and C. Rovelli. *The complete LQG propagator: I. Difficulties with the Barrett-Crane vertex*, *Phys. Rev.*, D76:104012, 2007, 0708.0883.
- [35] E. Alesci, E. Bianchi and C. Rovelli, “LQG propagator: III. The new vertex,” arXiv:0812.5018 [gr-qc].
- [36] B. Dittrich and J. P. Ryan, “Phase space descriptions for simplicial 4d geometries,” arXiv:0807.2806 [gr-qc].
- [37] A. Ashtekar, J. Lewandowski, D. Marolf, J. Mourao and T. Thiemann, “Coherent state transforms for spaces of connections,” *J. Funct. Anal.* **135**, 519 (1996) [arXiv:gr-qc/9412014].
- [38] B. Hall “The Segal-Bargmann coherent state’ transform for compact Lie groups” *Journal of functional analysis*, **122**, 103,1994.
- [39] B. Bahr and T. Thiemann, “Gauge-invariant coherent states for Loop Quantum Gravity I: Abelian gauge groups,” *Class. Quant. Grav.* **26** (2009) 045011 [arXiv:0709.4619 [gr-qc]].
- [40] L. Freidel and E. R. Livine, “Spin networks for non-compact groups,” *J. Math. Phys.* **44**, 1322 (2003) [arXiv:hep-th/0205268].